

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
24 January 2002 (24.01.2002)

PCT

(10) International Publication Number  
WO 02/06490 A1

- (51) International Patent Classification<sup>7</sup>: C12N 15/53, 15/82, 9/02, A01H 5/00, A01N 65/00
- (74) Agent: BASTIAN, Werner; c/o Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).
- (21) International Application Number: PCT/EP01/08085
- (22) International Filing Date: 12 July 2001 (12.07.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
0017275.9 13 July 2000 (13.07.2000) GB  
0022739.7 15 September 2000 (15.09.2000) GB
- (71) Applicants (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; Schwazwaldallee 215, CH-4058 Basel (CH). UNIVERSITÄT ZÜRICH [CH/CH]; Rämistrasse 71, CH-8006 Zürich (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): DUDLER, Robert [CH/CH]; Institut für Pflanzenbiologie, Universität Zürich, Zollikerstrasse 107, CH-8008 Zürich (CH). SCHAFFRATH, [DE/DE]; Institut für Biologie III, RWTH Aachen, Woringerweg 1, 52056 Aachen (DE). LAWTON, Kay, Ann [US/US]; Syngenta, 3054 Cornwallis Road, Research Triangle Park, NC 27709 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report
  - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/06490 A1

(54) Title: LIPOXYGENASE GENES, PROMOTERS, TRANSIT PEPTIDES AND PROTEINS THEREOF

(57) Abstract: This invention describes novel lipoxygenase genes and promoters, transit peptides and proteins derived therefrom. More specifically, this invention describes novel promoters that confer chemically inducible but not wound- or pathogen-inducible expression to an associated nucleotide sequence. The invention further describes peptides capable of targeting and associated protein to plastids and proteins having lipoxygenase activity which can be used to inhibit fungal mycotoxins. The invention also describes recombinant sequences containing such promoter sequences, and/or sequences encoding transit peptides and proteins according to the invention. The said recombinant DNA sequences may be used to create transgenic plants, but especially transgenic plants expressing a nucleotide sequence of interest in response to chemicals but not in response to wounding or pathogens.

Lipoxygenase Genes, Promoters, Transit Peptides and Proteins Thereof

The present invention relates to novel lipoxygenase genes and promoters, transit peptides and proteins derived therefrom. The present invention also relates the methods of using the novel lipoxygenase genes, promoters, transit peptides and proteins. The present invention also relates to isolated nucleic acid molecules encoding polypeptides having lipoxygenase activity and transit peptides. More specifically, this invention relates to isolated nucleic acid molecules encoding novel promoters that confer chemically inducible but not wound- or pathogen-inducible expression to an associated nucleotide sequence. Furthermore, the invention relates to peptides capable of targeting an associated protein to plastids. The invention also relates to proteins having lipoxygenase activity and to their use in inhibiting fungal mycotoxins. The invention further relates to recombinant nucleic acid molecules comprising nucleic acid molecules encoding the novel lipoxygenase genes, promoters or transit peptides. Also, the invention relates to host cells, plants or progeny thereof comprising the nucleic acid molecules or recombinant molecules described herein.

Plants are exposed to a variety of microbes during their life cycle, many of which are capable of causing disease. As a consequence plants have developed multiple defense strategies to avoid colonization. Certain treatments with chemical or biological agents can induce a normally susceptible plant to become systemically resistant to a subsequent inoculation with virulent pathogens. This phenomenon is known as systemic acquired resistance, or SAR.

In rice, for example, treatment with the chemical N-cyanomethyl-2-chloroisonicotinamid, a derivative of 2,6-dichloroisonicitnic acid (INA), has good resistance inducing activity against rice blast disease. Interestingly, treatment with N-cyanomethyl-2-chloroisonicotinamid, induced the enzyme lipoxygenase (LOX, linoleate:oxygen oxidoreductase, EC 1.13.11.12) (Seguchi et al. (1992, *Journal Pest. Sci.* 17, 107-113)), an enzyme known to be involved in plant defense against pathogens. Treatment with the rice blast fungus itself also induced this lipoxygenase. However, in modern agriculture, there is a desire to have a gene at hand, that is only induced by treatment with a chemical, but not by pathogens or wounding. Therefore, it is a major objective of the present invention to provide a lipoxygenase gene that is chemically induced, but not by pathogens or wounding.

It is another objective of the present invention, to provide the promoter and the transit peptide and protein encoded by such a lipoxygenase gene for use in agricultural biotechnology.

In agricultural biotechnology, plants can be modified according to one's needs. One way to accomplish this is by using modern genetic engineering techniques. For example, by introducing a gene of interest into a plant, the plant can be specifically modified to express a desirable phenotypic trait. For this, plants are transformed most commonly with a heterologous gene comprising a promoter region, a coding region and a termination region. When genetically engineering a heterologous gene for expression in plants, the selection of a promoter is often a critical factor. While it may be desirable to express certain genes constitutively, i.e. throughout the plant at all times and in most tissues and organs, other genes are more desirably expressed only in response to particular stimuli or confined to specific cells or tissues. Chemically inducible promoters have been previously described (see, for example EP A-332 104). However, these promoters are also induced by pathogens. There are however occasions where it is desirable to use a promoter that is chemically induced but not by pathogens or wounding. Therefore, it is a major objective of the present invention to provide such alternative promoters for expression of a nucleotide sequence of interest in plants. The invention also provides recombinant DNA molecules, expression vectors and transgenic plants comprising the promoters of the present invention. When genes of interest are introduced into plants, they are most commonly expressed in the cytoplasm. Alternatively, one might wish to express those genes in other compartments of the cell. This can be accomplished, for example, by introducing the gene of interest into the plastid genome instead of the nuclear genome. However, currently, plastid transformation is not a routine procedure for all of the agriculturally important crops. Another possible way to express a protein of interest in plastids is to add a DNA sequence encoding a transit peptide to the 5'-end of the DNA sequence encoding a protein of interest and to express this DNA sequence from the nuclear genome. Transit peptides are peptides that are capable of targeting an associated protein to plastids. It is thus another objective of the present invention to provide such transit peptides. The invention also provides recombinant DNA molecules, expression vectors and transgenic plants comprising the transit peptides of the present invention. The transit peptides can be used in completely heterologous constructs or together with the promoter or coding region they are naturally associated with. The present invention also provides recombinant DNA molecules, expression vectors and transgenic plants comprising the transit peptides of the present invention.

In agricultural biotechnology not only the choice of the promoter is of importance, but also the choice of the associated DNA encoding a desirable phenotypic trait. A particularly desirable phenotypic trait is the lipoxygenase protein of the present invention. The invention thus provides recombinant DNA molecules, expression vectors and transgenic plants comprising the lipoxygenase protein of the present invention.

The present invention thus provides:

an isolated nucleic acid molecule capable of driving chemically inducible but not wound- or pathogen-inducible expression of an associated nucleotide sequence in particular, wherein said isolated nucleic acid molecule

- is a component of the *Pst*I/*Pst*I fragment of about 4.5 kb in length from plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524
- is a component of the nucleotide sequence depicted in SEQ ID NO:17
- is depicted in SEQ ID NO:18
- is depicted in SEQ ID NO:19
- comprises the nucleotide sequence depicted in SEQ ID NO:1
- comprises the nucleotide sequence depicted in SEQ ID NO:2
- comprises nt 1 to nt 1358 of the nucleotide sequence depicted in SEQ ID NO:2
- comprises the nucleotide sequence depicted in SEQ ID NO:3
- comprises nt 1702 to nt 2104 of SEQ ID NO:2 and/or nt 1 to nt 97 of SEQ ID NO:3 and/or nt 367 to nt 1283 of SEQ ID NO:3 of SEQ ID NO:3
- comprises a combination of any one of the nucleotide sequences or portions thereof depicted in SEQ ID NO:1, SEQ ID NO:2 and SEQ ID NO:3
- hybridizes under stringent conditions to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or to the 4.5 kb *Pst*I fragment of plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524, wherein said nucleic acid molecule is capable of driving chemically inducible but not wound- or pathogen-inducible expression of an associated nucleotide sequence
- comprises a consecutive stretch of at least 50 nt, preferably of about 500 bases, particularly of between about 1000 bases and about 1500 bases, more particularly of about 2000 bases and most particularly of between about 3000 bases and about 4500 bases in length of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or of the 4.5 kb *Pst*I fragment of plasmid pBSK+LOX4A which

has been deposited under accession no DSM 13524, wherein said isolated nucleic acid molecule is capable of driving chemically inducible but not wound- or pathogen-inducible expression of an associated nucleotide sequence, in particular, wherein said consecutive stretch of at least 50 nt has at least 70%, preferably 80%, more preferably 90% and most preferably 95% sequence identity with a consecutive stretch of corresponding length of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or the 4.5 kb *Pst*I fragment of plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524

- wherein the chemical inducer capable of inducing said nucleic acid molecule is selected from the group consisting of BTH (benzo(1,2,3)thiadiazole-7-carbothioic acid S-methyl ester), INA (2,6-dichloroisonicotinic acid) and probenazole
- wherein the chemical inducer capable of inducing said nucleic acid molecule is jasmonic acid

Further provided are recombinant nucleic acid molecules comprising a nucleic acid molecule according to the invention operably linked to a nucleotide sequence of interest in particular, wherein

- the nucleotide sequence of interest comprises a protein, polypeptide or peptide coding sequence
- the coding sequence comprises at its 5'-end a nucleotide sequence encoding the amino acid sequence depicted in SEQ ID NO:6
- the coding sequence encodes a desirable phenotypic trait
- the coding sequence encodes a selectable or screenable marker gene
- the coding sequence encodes a protein conferring antibiotic resistance, virus resistance, insect resistance, disease resistance, or resistance to other pests, herbicide tolerance, improved nutritional value, improved performance in an industrial process or altered reproductive capability
- the coding sequence encodes commercially valuable enzymes or metabolites in the plant
- the coding sequence is in antisense orientation

Further provided are isolated nucleic acid molecules expression vectors comprising an isolated nucleic acid molecule or a recombinant nucleic acid molecule of the invention as well as host cells stably transformed with a isolated nucleic acid molecule or a recombinant nucleic acid molecule according to the invention in particular, wherein

- the host cell is a bacterium

- the host cell is a plant cell
- the host cell is a plant cell selected from the group consisting of rice, maize, wheat, barley, rye, sweet potato, sweet corn, bean, pea, chicory, lettuce, cabbage, cauliflower, broccoli, turnip, radish, spinach, asparagus, onion, garlic, pepper, celery, squash, pumpkin, hemp, zucchini, apple, pear, quince, melon, plum, cherry, peach, nectarine, apricot, strawberry, grape, raspberry, blackberry, pineapple, avocado, papaya, mango, banana, soybean, tomato, sorghum, sugarcane, sugar-beet, sunflower, rapeseed, clover, tobacco, carrot, cotton, alfalfa, potato, eggplant, cucumber, *Arabidopsis thaliana*, and woody plants such as coniferous and deciduous trees, but particularly rice, maize, wheat, barley, cabbage, cauliflower, pepper, squash, melon, soybean, tomato, sugar-beet, sunflower or cotton, rice, maize, wheat, *Sorghum bicolor*, orchardgrass, sugar beet and soybean cells
- the host cell is a plant cell from a dicotyledonous plant
- the host cell is a plant cell from a dicotyledonous plant selected from the group consisting of soybean, cotton, tobacco, sugar beet and oilseed rape
- the host cell is a plant cell from a monocotyledonous plant
- the host cell is a plant cell from a monocotyledonous plant selected from the group consisting of maize, wheat, sorghum, rye, oats, turf grass, rice, and barley.

Further provided are plants and the progeny thereof stably transformed with a nucleic acid molecule or a recombinant nucleic acid molecule according to the invention. In particular, wherein said plant is selected from the group consisting of maize, wheat, sorghum, rye, oats, turf grass, rice, barley, soybean, cotton, tobacco, sugar beet and oilseed rape. Further provided are seeds from the transformed plants and progeny thereof.

In addition, use of the isolated nucleic acid molecule of the invention to express a nucleotide sequence of interest is provided.

The present invention further discloses

- the use of the isolated nucleic acid molecule according to the invention to express a nucleotide sequence of interest
- a method of producing an isolated nucleic acid molecule according to the invention, wherein the nucleic acid molecule is produced by a polymerase chain reaction wherein at least one oligonucleotide used comprises a sequence of nucleotides which represents a

•consecutive stretch of 15 or more base pairs of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19.

The invention also provides isolated nucleic acid molecules encoding the amino acid sequence depicted in SEQ ID NO:6, wherein said amino acid sequence is capable of targeting an associated protein to plastids in particular, wherein

- said nucleotide sequence is the sequence depicted in SEQ ID NO:4
- said nucleotide sequence hybridizes under stringent conditions to SEQ ID NO:4 in particular, wherein said sequence has 70%, preferably 80%, more preferably 90% sequence identity with the nucleotide sequence of SEQ ID NO:4 and the encoded peptide is capable of targeting an associated protein to plastids

Further are provided polypeptides or peptides encoded by the isolated nucleic acid molecules described above as well as the use of said polypeptides or peptides to target an associated protein of interest to plastids.

In addition, the invention provides isolated nucleic acid molecules which hybridize under stringent conditions to SEQ ID NO:5, and wherein the protein encoded by said nucleic acid molecule has at least 65%, preferably 75% more preferably 85 % and most preferably 95% amino acid sequence identity with the amino acid sequence depicted in SEQ ID NO:7 and encodes a protein with lipxygenase activity.

The invention further provides nucleic acid molecules as mentioned hereinbefore, wherein said nucleic acid molecules encode the protein depicted in SEQ ID NO:7. Further are provided proteins encoded by said nucleic acid molecules described hereinbefore, in particular, SEQ ID NO:7 or portions of the proteins or polypeptides having lipxygenase activity.

The invention further discloses the use of the protein as mentioned hereinbefore to inhibit fungal mycotoxins, in particular aflatoxins. The invention further provides methods of increasing plant disease resistance or inhibiting fungal mycotoxins by expressing the isolated nucleic acid molecules of the present invention that encode lipxygenase activity in transformed plants.

Further provided are recombinant nucleic acid molecule comprising the nucleic acid molecules as described above, host cells stably transformed therewith, in particular wherein said host cell is a plant cell and plants and the progeny thereof stably transformed with a recombinant nucleic acid molecule as described above.

In order to ensure a clear and consistent understanding of the specification and the claims, the following definitions are provided:

DNA shuffling: DNA shuffling is a method to rapidly, easily and efficiently introduce rearrangements, preferably randomly, in a DNA molecule or to generate exchanges of DNA sequences between two or more DNA molecules, preferably randomly. The DNA molecule resulting from DNA shuffling is a shuffled DNA molecule that is a non-naturally occurring DNA molecule derived from at least one template DNA molecule.

Expression: refers to the transcription and/or translation of an endogenous gene or a transgene in plants. In the case of antisense constructs, for example, expression may refer to the transcription of the antisense DNA only.

Functionally equivalent sequence: refers to a DNA sequence which has promoter activity substantially similar to the rice lipoxygenase gene promoter or parts thereof and which under stringent hybridizing conditions hybridizes with the said promoter sequences.

Gene: refers to a coding sequence and associated regulatory sequence wherein the coding sequence is transcribed into RNA such as mRNA, rRNA, tRNA, snRNA, sense RNA or antisense RNA. Examples of regulatory sequences are promoter sequences, 5'- and 3'- untranslated sequences and termination sequences. Further elements that may be present are, for example, introns.

Gene of interest: refers to any gene which, when transferred to a plant, confers upon the plant a desired characteristic such as antibiotic resistance, virus resistance, insect resistance, disease resistance, or resistance to other pests, herbicide tolerance, improved nutritional value, improved performance in an industrial process or altered reproductive capability. The "gene of interest" may also be one that is transferred to plants for the production of commercially valuable enzymes or metabolites in the plant.

Heterologous as used herein means of different natural or of synthetic origin. For example, if a host cell is transformed with a nucleic acid sequence that does not occur in the untransformed host cell, that nucleic acid sequence is said to be heterologous with respect to the host cell. The transforming nucleic acid may comprise a heterologous promoter, heterologous coding sequence, or heterologous termination sequence. Alternatively, the transforming nucleic acid may be completely heterologous or may comprise any possible combination of heterologous and endogenous nucleic acid sequences.

Leader region: region in a gene between transcription start site and translation start site.

LOX: lipoxygenase.

Marker gene: refers to a gene encoding a selectable or screenable trait.



nt: nucleotide, and are naturally occurring or synthetic nucleotides.

Nucleic acid molecule: is any single or double stranded polynucleotide that is commonly either DNA or RNA, and can comprise naturally occurring or synthetic nucleotides.

Operably linked to/associated with: a regulatory DNA sequence is said to be "operably linked to" or "associated with" a DNA sequence that codes for an RNA or a protein if the two sequences are situated such that the regulatory DNA sequence affects expression of the coding DNA sequence.

Plant: refers to any plant, particularly to seed plants.

Plant cell: structural and physiological unit of the plant, comprising a protoplast and a cell wall. The plant cell may be in form of an isolated single cell or a cultured cell, or as a part of higher organized unit such as, for example, a plant tissue, or a plant organ.

Plant material: refers to leaves, stems, roots, flowers or flower parts, fruits, pollen, pollen tubes, ovules, embryo sacs, egg cells, zygotes, embryos, seeds, cuttings, cell or tissue cultures, or any other part or product of a plant

Polynucleotide: any single-stranded homo-or heteropolymer of at least about ten nucleotides connected by phosphodiester linkages between (usually) the 3' position of the glycoside moiety of one nucleotide and the 5' position on the glycoside moiety of the adjacent nucleotide, or any double-stranded molecule comprised of two such single-stranded molecules held together by hydrogen bonds.

Promoter: refers to a DNA sequence that initiates transcription of an associated DNA sequence. The promoter region may also include elements that act as regulators of gene expression such as activators, enhancers, and/or repressors and may include all or part of the 5' non-translated region.

Protein, Polypeptide or peptide: are used herein interchangeably and are amino acid residues connected by peptide linkages.

Recombinant DNA molecule: a combination of DNA sequences that are joined together using recombinant DNA technology.

Recombinant DNA technology: procedures used to join together DNA sequences as described, for example, in Sambrook et al., 1989, Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.

Screenable marker gene: refers to a gene whose expression does not confer a selective advantage to a transformed cell, but whose expression makes the transformed cell phenotypically distinct from untransformed cells.

Selectable marker gene: refers to a gene whose expression in a plant cell gives the cell a selective advantage. The selective advantage possessed by the cells transformed with the selectable marker gene may be due to their ability to grow in the presence of a negative selective agent, such as an antibiotic or a herbicide, compared to the growth of non-transformed cells. The selective advantage possessed by the transformed cells, compared to non-transformed cells, may also be due to their enhanced or novel capacity to utilize an added compound as a nutrient, growth factor or energy source. Selectable marker gene also refers to a gene or a combination of genes whose expression in a plant cell in the presence of the selective agent, compared to the absence of the selective agent, has a positive effect on the transformed plant cell and a negative effect on the un-transformed plant cell, for example with respect to growth, and thus gives the transformed plant cell a positive selective advantage.

Sequence identity: the percentage of sequence identity is determined using computer programs that are based on dynamic programming algorithms. Computer programs that are preferred within the scope of the present invention include the BLAST (Basic Local Alignment Search Tool) search programs designed to explore all of the available sequence databases regardless of whether the query is protein or DNA. Version BLAST 2.0 (Gapped BLAST) of this search tool has been made publicly available on the Internet (currently <http://www.ncbi.nlm.nih.gov/BLAST/>). It uses a heuristic algorithm, which seeks local as opposed to global alignments and is therefore able to detect relationships among sequences, which share only isolated regions. The scores assigned in a BLAST search have a well-defined statistical interpretation. Said programs are preferably run with optional parameters set to the default values.

Transformation: refers to the introduction of a nucleic acid into a cell. In particular, it refers to the stable integration of a DNA molecule into the genome of an organism of interest.

The present invention relates to lipoxygenase genes, and to promoters, transit peptides and proteins derived therefrom. Preferred are lipoxygenase genes that are chemically induced, but not by pathogens or wounding. In particular, said lipoxygenase genes are from rice. Such lipoxygenase genes, or portions or fragments therefrom, can be obtained, for example, by a PCR-based strategy. For this, known lipoxygenase coding sequences, for example, from rice (Peng et al. (1994) *J. Biol. Chem.* 269, 3755-3761; Ohta et al. (1992) *Eur. J. Biochem.* 206, 331-336) and from wheat (Görlach et al. (1996) *Plant Cell.* 8, 629-643) are aligned to identify conserved regions using computer programs known in the art.

By this method, several conserved regions are identified, one of which is near the C-terminus and contains the amino acid sequence HAAVNFG that is invariant in all three sequences. Then, total RNA or polyA<sup>+</sup> RNA is isolated from untreated control leaves and from leaves sprayed with a 100 ppm INA solution and harvested 24 and 48 hours after treatment.

The RNA samples are used as templates for RT-PCR using the degenerate oligonucleotide 5'-CAYGCNGTNAANTTYGG-3' (SEQ ID NO:8), which corresponds to the HAAVNFG amino acid sequence motif in the C-terminal region of the rice RLL2 lipxygenase (Peng et al. (1994) *J. Biol. Chem.* 269, 3755-3761), as the forward primer and an anchored oligo-dT primer as the reverse primer (5'-AATGCTTTTTTTTTTTTTTV-3', SEQ ID NO:9). When this method is performed with total RNA or polyA<sup>+</sup> RNA from rice, a PCR product of approximately 600 bp arises on ethidiumbromide stained agarose gels only in the INA-treated sample but not in the control. Those of skill in the art know that the size of the band can be smaller or larger, depending on the organism from which the RNA is isolated. The obtained band can be cloned and sequenced and used as a probe to screen cDNA or genomic libraries to obtain full-length lipxygenase cDNA or genomic clones by methods known in the art. Upon screening a rice cDNA library constructed from INA-treated leaves, a full-length rice lipxygenase cDNA clone of 3018 bp in length is obtained (SEQ ID NO:5). This cDNA clone, designated RCI-1 (rice chemically induced cDNA 1), contains an open reading frame of 2766 bp (from base 48 to base 2816 of SEQ ID NO:5) encoding a protein of 922 amino acid residues with a predicted Mr of 105 kDa (SEQ ID NO:7). To those with skill in the art it is known that the obtained cDNA clone can be larger or smaller, depending on whether the clone is full-length or not, on the length of the 5' and 3' untranslated region, and on the organism from which the library is constructed.

When the RCI-1 cDNA is used as a probe in Northern blot analyses with RNA from chemically treated leaves, such as leaves treated with INA, BTH, probenazole or jasmonic acid, a strong hybridization signal is observed, indicating the accumulation of RCI-1 mRNA. No such mRNA accumulation is observed when RNA from wounded or pathogen-treated leaves is used. This is surprising, as wounding is known to increase endogenous levels of jasmonic acid in rice and induces increased systemic protection against rice blast infection (Schweizer et al. (1998) *Plant J.* 14, 475-481; Schweizer et al. (1997) *Plant Physiol.* 114, 79-88). The lipxygenase of the present invention however is not induced by pathogens such as the rice blast fungus *Magnaporthe grisea* nor the bacterial pathogen *Pseudomonas syringae* pv. *syringae*. This indicates that the promoter region of the corresponding gene

must contain regulatory elements that confer a chemically, but not wound- or pathogen inducible expression pattern to an associated coding sequence.

The protein encoded by the RCI-1 cDNA is most similar to the barley LOX2:Hv:1 (60% identity and 68% similarity). Sequence identity (similarity) at the amino acid level are 43% (52%) for the rice lipoxygenase L-2 predominately found in kernels and seedlings (Ohta et al. (1992) *Eur. J. Biochem.* 206, 331-336) and 50% (58%) for the *Magnaporthe grisea*-induced rice lipoxygenase RLL2 (Peng et al. (1994) *J. Biol. Chem.* 269, 3755-3761). DNA sequences embraced by the present invention are those that hybridize to the RCI-1 cDNA clone (SEQ ID NO:5) under stringent conditions and whose coding sequences have at least 65%, preferably 75%, more preferably 85 % and most preferably 95% amino acid sequence identity to the protein depicted in SEQ ID NO:7 and encode a protein with lipoxygenase activity.

The lipoxygenase cDNA of the present invention can be expressed in *E. coli* or in any other expression system suitable to express eukaryotic sequences by methods known in the art. The expressed protein is then analyzed and, optionally, purified. All these methods are known to a person skilled in the art. When an extract of *E. coli* cells expressing a cDNA of the present invention is analyzed, increased LOX activity using linoleic acid as a substrate is detected, while control extracts of *E. coli* without expression construct or containing the empty vector do not have detectable LOX activity. Maximal activity is observed around pH 8 to 9, indicating that RCI-1 must be classified as a type 1 LOX (Siedow (1991) *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42, 145-188). However, it should be noted that recently a second classification based on the presence of a plastomic transit peptide was introduced (Shibata et al. (1994) *Plant Mol. Biol. Rep.* 12, 41-42). According to this scheme, RCI-1 must be classified as a type 2 LOX.

When the reaction products of the recombinant lipoxygenase of the present invention are analyzed by HPLC (Bohland et al. (1997) *Plant Physiol.* 114, 679-685), (13S)-hydroperoxy-(9Z, 11E, 15Z)-octadecatrienoic acid (13-HPOD) is the predominant product, irrespective of whether linoleic acid or linolenic acid serves as a substrate for the enzyme. (9S)-hydroperoxy-(10E, 12Z, 15Z) octadecatrienoic acid (9-HPOD) is only detected in minor amounts. Reaction products derived from 13-HPOD have been reported to act as antimicrobial substances against *Magnaporthe grisea* (Shimura et al., (1981) *Agric Biol Chem* 45: 1431-1435; Shimura et al (1983) *Agric Biol Chem* 47: 1983-1989). This indicates that the lipoxygenase of the present invention, in particular RCI-1 LOX, is involved in the

generation of fatty acid derivatives that may act as signals or exhibit direct antimicrobial activity (reviewed in Slusarenko, A. J. (1996) The role of lipoxygenase in resistance to infection. In *Lipoxygenase and Lipoxygenase Pathway Enzymes* (Piazza, G. J., ed) pp. 176-197. American Oil Chemists Society Press, Champaign, Illinois). In one particular embodiment, the lipoxygenase of the present invention is suited to eliminate or substantially reduce the activity of fungal mycotoxins, which include, but are not limited to aflatoxins and their precursor sterigmatocystin, citrinin, fungal tremorgens, lupinosis, ochratoxins, patulin, rubratoxins, sporidesmin, stachybotryotoxins, trichothecens and zearalenone, but particularly aflatoxin and sterigmatocystin.

An alignment of the amino acid sequence of the lipoxygenase protein of the present invention with the amino acid sequence of other lipoxygenases such as barley LoxA (GenBank accession number L35931) or rice L-2 (Swissprot accession number P29250) shows that the lipoxygenase of the present invention has a N-terminal extension of between 30 and 50 amino acid residues. In the case of the rice lipoxygenase RCI-1 that extension is about 47 amino acids in length. This N-terminal extension clearly separates this class of LOX species from another class that is predominately found in kernels and seedlings and that includes LoxA, LoxB, and LoxC from barley (van Mechelen et al. (1999) *Plant Mol. Biol.* 39, 1283-1298), and LOX L-2 from rice (Ohta et al (1992) *Eur. J. Biochem.* 206, 331-336). When parts or all of this N-terminal extension is fused to the N-terminal region of a reporter gene, the reporter gene is targeted to plastids, in particular to chloroplasts. For example, when a chimeric gene is constructed with the first 158 bp of the *RCI-1* cDNA (SEQ ID NO:4) fused to the 5' end of the coding sequence of the green fluorescent protein (GFP), a modified GFP is obtained which contains at its N-terminus the first 37 amino acids of RCI-1 (SEQ ID NO:6). When said construct is introduced into Arabidopsis tissue, for example, by *Agrobacterium* based transformation system, a strict congruence of GFP fluorescence and chlorophyll autofluorescence is observed, indicating that the fusion protein is localized in the chloroplasts. Thus, the N-terminal extensions of the lipoxygenase proteins of the present invention function as transit peptides to transfer associated proteins to plastids, particularly to chloroplasts.

In addition, the present invention also provides promoters capable of conferring chemically inducible, but not wound- or pathogen-inducible expression to an associated nucleotide sequence of interest. Preferred are promoter sequences obtainable from the rice lipoxygenase gene RCI-1. Nucleotide sequences comprising functional and/or structural

equivalents thereof are also embraced by the invention. The present invention thus relates to nucleotide sequences that function as promoters of transcription of associated nucleotide sequences. The promoter region may also include elements that act as regulators of gene expression such as activators, enhancers, and/or repressors and may include the 5' non-translated leader sequence of the transcribed mRNA and/or introns and, optionally, exons. Chemically inducible, but not wound- or pathogen inducible expression means that the nucleotide sequence of interest is preferentially expressed when a chemical compound according to the invention is applied, but not upon wounding or exposure to pathogens. Thus, the nucleotide sequence according to the invention is useful for chemically inducible, but not wound- or pathogen inducible expression of an associated nucleotide sequence of interest, which preferably is a coding sequence. It is known to the skilled artisan that the associated coding sequence of interest can be expressed in sense or in antisense orientation. Further, the coding sequence of interest may be of heterologous or homologous origin with respect to the plant to be transformed. In case of a homologous coding sequence, the nucleotide sequence according to the invention is useful for ectopic expression of said sequence. In one particular embodiment of the invention expression of the coding sequence of interest under control of a nucleotide sequence according to the invention suppresses its own expression and that of the original copy of the gene by a process called co-suppression.

The promoters of the present invention can be obtained, for example, from rice genomic DNA by probing a rice genomic library with a cDNA according to the invention using methods known in the art. It is obvious to a person skilled in the art that genomic DNA from any other organism, particularly from plants, can be used to obtain a lipoxygenase promoter from any organism of interest. This genomic DNA is then sequenced and aligned to the cDNA sequence. Basically, all nucleotide sequences upstream of the start codon are considered to be part of the lipoxygenase promoter region. In addition, introns and, optionally, exons can be added to this region to form a functional promoter that confers chemically inducible, but not wound- or pathogen inducible expression to an associated coding region.

In a preferred embodiment of the invention, the lipoxygenase promoter is a component of the *Pst*I/*Pst*I fragment of about 4.5 kb in length from plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524. SEQ ID NO:17 depicts the nucleotide sequence of the about 4.5 kb *Pst*I/*Pst*I fragment from plasmid pBSK+LOX4A. Other preferred

embodiments of the invention are the nucleotide sequences depicted in SEQ ID Nos:18, 19, 1, 2 and 3, which are components of the 4.5 kb *Pst*I/*Pst*I fragment as mentioned hereinbefore. Another preferred embodiment of the invention comprises nt 1 to nt 1358 of the nucleotide sequence depicted in SEQ ID NO:2.

SEQ ID NO:1 comprises the 5'-end of the 4.5 kb *Pst*I/*Pst*I fragment. This nucleotide sequence is 358 nucleotides in length and contains at its 5' end in position 1 to 6 the *Pst*I-site. The region between SEQ ID NO:1 and SEQ ID NO:2 of the 4.5 kb *Pst*I/*Pst*I fragment is between about 240 and 440 bp in length. The central region of the 4.5 kb *Pst*I/*Pst*I fragment is shown in SEQ ID NO:2 and is 2104 bp in length. It contains the putative TATA box (position 1261 to 1266 SEQ ID NO:2), the putative start codon (position 1359 to 1361 of SEQ ID NO:2), as well as the 5'-untranslated region and nucleotide sequences upstream of the putative TATA box. Comparison of the genomic DNA (SEQ ID NO:2) and the cDNA (SEQ ID NO:5) shows that the sequences located at position 1312 to 1701 of SEQ ID NO:2 comprise all or part of exon 1, and the sequences located at position 1702 to 2104 of SEQ ID NO:2 are the 5' part of intron 1. The region between SEQ ID NO:2 and SEQ ID NO:3 of the 4.5 kb *Pst*I/*Pst*I fragment is between about 85 and 285 bp in length. The 3' end of the 4.5 kb *Pst*I/*Pst*I fragment is shown in SEQ ID NO:3. This sequence depicts a nucleotide sequence of 1516 bp in length. It contains, in a 5' to 3' direction, the 3' end of intron 1 (position 1 to 97 of SEQ ID NO:3) followed by exon 2 (position 98 to 366 of SEQ ID NO:3), intron 2 (position 367 to 1283 of SEQ ID NO:3) and part of exon 3 (position 1284 to 1516 of SEQ ID NO:3). The *Pst*I site is located at position 1511 to 1516.

Based on the sequence information given in SEQ ID NOs:1 to 3, the DNA sequences of the invention can be obtained, for example, by PCR using plasmid pBSK+LOX4A or genomic DNA from rice or any other organism of interest as template. The person skilled in the art knows how to arrive at such sequences using methods known in the art. These sequences then can be fused to reporter genes to demonstrate promoter activity. For example, chimeric genes can be constructed that include part of the 5' regulatory sequence of the RCI-1 gene fused to the GFP coding sequence. To this end, pBSK+LOX4A (see Example 9) can be used as template for the polymerase chain reaction (PCR). Gene-specific primers can be designed to amplify the 5' promoter region of the gene. Using combinations of, for example, the reverse primer R1 (SEQ ID NO:12) with forward primers F1 (SEQ ID NO:13) and F2 (SEQ ID NO:14) the regulatory sequences that are ~1.2 kb and ~2 kb upstream of

the initiating methionine are isolated. The nucleotide sequence of the PCR fragment amplified with forward primer F1 and reverse primer R1 is shown in SEQ ID NO:18, and the nucleotide sequence of the PCR fragment amplified with forward primer F2 and reverse primer R1 is shown in SEQ ID NO:19. For ease of cloning the primers consist, for example, of gene specific sequences and attB recombination sites for the GATEWAY™ cloning technology (Life Technologies, GIBCO BRL, Rockville, MD USA). As reverse primer, primer R1 can be used, which has the following sequence: 5'-

CAAGAAAGCTGGGTTGACAAATTAAGTTGTCAGTGTG-3' (SEQ ID NO:12). The gene specific sequence of reverse primer R1 is underlined (corresponds to position 1356 to 1334 of SEQ ID NO:2), the attB recombination sequence is denoted in italics. Examples for forward primers are the primers F1 and F2. Forward primer F1 has the following sequence: 5'-CAAAAAAGCAGGCTTGTAACATCCTACTCCTATTGTG-3' (SEQ ID NO:13). The gene specific sequence of forward primer F1 is underlined (corresponds to bases 159 to 181 of SEQ ID NO:2), the attB recombination sequence is denoted in italics. F1 in combination with R1 amplifies a fragment of ~1.2 kb. Forward primer F2 has the following sequence: 5'-CAAAAAAGCAGGCTCCCGTCTTTATCTACTC-3' (SEQ ID NO:14). The gene specific sequence of forward primer F2 is underlined (corresponds to bases 31 to 48 of SEQ ID NO:1), the attB recombination sequence is denoted in italics. Primer F2 in combination with primer R1 amplifies a fragment of ~2 kb.

Using a nested PCR strategy the regulatory sequence can be amplified first with primers F1+R1 or F2+R1 followed by a second PCR with primer attB1 (5'-GGGGACAAGTTTGTACAAAAAAGCAGGCT-3', SEQ ID NO:15) and primer attB2 (5'-GGGGACCACTTTGTACAAGAAAGCTGGGT-3', SEQ ID NO:16). Optimal annealing temperatures can be determined using a gradient thermocycler (DNA Engine, MJ Research, Inc. Waltham, MA USA) and the following PCR conditions with gene-specific primers F1+R1 or F2+R1: [(94°C:10sec):(94°C:10sec/45°C to 70°C gradient:10sec/72°C:10sec)X15]. Following PCR the products can be visualized by gel electrophoresis, and DNA from the reaction with the highest T<sub>m</sub> giving visible product can be selected for amplification with the attB1+attB2 primers. In the subsequent PCR amplification, the following PCR conditions can be used: [(94°C:10sec):(94°C:10sec/50°C to 70°C gradient 10sec/72°C:10sec)X15]. The resulting PCR product are then flanked by attB recombination sites which can be used to generate Entry Clones in pENTR via the BP reaction according to manufacturers protocol (see: Instruction Manual of GATEWAY™ Cloning Technology, GIBCO BRL, Rockville, MD



USA, <http://www.lifetech.com/>). The resulting plasmids contain ~1.2 kb and ~2 kb 5' of the *RCI-1* initiation codon and are referred to as pENTR+LOXp1.2, pENTR+LOXp2.

The regulatory/promoter sequence is then fused to the mGFP-5 reporter gene (MRC Laboratory of Molecular Biology, Cambridge, England) by recombination using GATEWAY™ Technology according to manufacturers protocol as described in the Instruction Manual (GATEWAY™ Cloning Technology, GIBCO BRL, Rockville, MD <http://www.lifetech.com/>). Briefly, according to this protocol the promoter fragment in the entry vector is recombined via the LR reaction with a binary *Agrobacterium* destination vector containing the mGFP-5 coding region that has an attR site 5' to the GFP reporter. The orientation of the inserted fragment is maintained by the att sequences and the final construct is verified by sequencing. The construct is designated pLOXp1.2 promoter::GFP or pLOXp2promoter::GFP and can be transformed into *Agrobacterium tumefaciens* strains by electroporation. Any other binary vector can also be modified to accommodate promoter fragments of the invention to drive expression of an associated reporter gene. The skilled artisan knows how to construct such vectors starting from commercially available binary vectors such as, for example pGPTV-BLEO (ATCC number 77392), pBI 121 (Clontech, Palo Alto, California), or pCambia 1302 (Cambia, Canberra, Australia).

Transgenic plants are then produced using, for example, *Agrobacterium*-mediated transformation techniques. Expression of the gene fusion protein can be monitored in transformants by confocal imaging using a Leica-TCS confocal laser scanning microscope and a PLAPO x100 oil immersion objective (Leica Microsystems, Heidelberg, Germany) with the following filter settings: excitation 476/488 nm; GFP-emission 515-552 nm, chlorophyll-emission 673-695 nm. GFP fluorescence and chlorophyll autofluorescence are recorded simultaneously using independent 2-channel-detection.

Confocal imaging of leaves from transgenic rice plants expressing the pRCI promoter::GFP construct can be carried out to assay promoter activity in response to abiotic and biotic inducers.

It is apparent to the skilled artisan that, based on the nucleotide sequences shown in SEQ ID NO:1 to 3, any primer combination of interest can be chosen to PCR amplify DNA fragments of various lengths that can be used according to the invention. Thus, any region of interest can be amplified from SEQ ID NOs:1 to 3. For example, primers can be designed to specifically amplify intron 1 or intron 2 or the 5' upstream region. The 5' upstream region is defined herein as the region between the putative TATA box and the putative start codon

of the lipoxygenase protein. The skilled artisan also will consider to combine intron 1 and/or intron 2 with various parts of SEQ ID NOs:1, 2 and/or 3, such as to arrive at an DNA molecule comprising nt 1702 to nt 2104 of SEQ ID NO:2 and/or nt 1 to nt 97 of SEQ ID NO:3 and/or nt 367 to nt 1283 of SEQ ID NO:3 of SEQ ID NO:3.

Further, it might also be desirable to combine any of these sequences with the 3' untranslated region of the lipoxygenase cDNA sequence (position 2817 to 3018 of SEQ ID NO:5).

The invention thus includes fragments derived from the rice RCI-1 lipoxygenase gene that function according to the invention, i.e. are capable of conferring chemically induced but not wound- or pathogen induced expression of an associated nucleotide sequence.

This can be tested by generating such promoter fragments, fusing them to a selectable or screenable marker gene and assaying the fusion constructs for retention of promoter activity in transient expression assays with protoplasts or in stably transformed plants. Such assays are within the skill of the ordinary artisan. Preferred nucleic acid molecule fragments of the invention are of at least about 500 bases, particularly of between about 1000 bases and about 1500 bases, more particularly of about 2000 bases and most particularly of between about 3000 bases and about 4500 bases in length.

It is also clear to the skilled artisan that mutations, insertions, deletions and/or substitutions of one or more nucleotides can be introduced into the nucleotide sequences of SEQ ID NOs:1, 2 and 3 or longer or shorter fragments derived from the sequence information thereof using methods known in the art. In addition, an unmodified or modified nucleotide sequence of the present invention can be varied by shuffling the sequence of the invention. To test for a function of variant nucleotide sequences according to the invention, the sequence of interest is operably linked to a selectable or screenable marker gene and expression of the marker gene is tested in transient expression assays with protoplasts or in stably transformed plants. It is known to the skilled artisan that nucleotide sequences capable of driving expression of an associated nucleotide sequence are build in a modular way. Accordingly, expression levels from shorter nucleic acid molecule fragments may be different than the one from the longest fragment and may be different from each other. For example, deletion of a down-regulating upstream element will lead to an increase in the expression levels of the associated nucleotide sequence while deletion of an up-regulating element will decrease the expression levels of the associated nucleotide sequence.

Another way of identifying promoter elements necessary for regulated expression of an associated nucleotide sequence is the so-called linker-scanning analysis. Linker-scanning

mutagenesis allows for the identification of short defined motifs whose mutation alters the promoter activity. Accordingly, a set of linker-scanning mutant promoters fused to the coding sequence of the GUS reporter gene or another marker gene can be constructed using methods known in the art. These constructs are then transformed into *Arabidopsis*, for example, and GUS activity is assayed in several independent transgenic lines. The effect of each mutation on promoter activity is then compared to an equivalent number of transgenic lines containing an unmutated rice lipoxygenase gene promoter. It is expected, that when a motif is mutated that is involved in chemically, but not wound or pathogen-inducible expression, that the level of expression of the reporter gene is modified. If, for example, a higher average induction of GUS activity by a chemical inducer is detected than the one from the control construct most likely a negative regulatory element had been mutated in this construct. If, on the other hand, a complete loss of inducibility of GUS activity by a chemical regulator according to the invention is observed, most likely a positive regulatory element necessary for chemical induction has been mutated. In a next step, particularly in the case of the putative positive regulatory element, the wild-type sequences corresponding to the mutated fragments are fused to a minimal promoter and the newly created promoter is tested for the ability to confer regulated expression to an associated marker gene.

Embraced by the present invention are also functional equivalents of the RCI-1 promoters of the present invention, i.e. nucleotide sequences that hybridize under stringent conditions to any one of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or to the 4.5 *Pst*I/*Pst*I fragment of plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524. A stringent hybridization is performed at a temperature of 65°C, preferably 60°C and most preferably 55°C in double strength (2X) citrate buffered saline (SSC) containing 0.1% SDS followed by rinsing of the support at the same temperature but with a buffer having a reduced SSC concentration. Such reduced concentration buffers are typically one tenth strength SSC (0.1X SSC) containing 0.1% SDS, preferably 0.2X SSC containing 0.1% SDS and most preferably half strength SSC (0.5X SSC) containing 0.1% SDS. In fact, functional equivalents to all or part of the RCI-1 lipoxygenase promoter from other organisms can be found by hybridizing any one of SEQ ID NO:1, SEQ ID NO:2, or SEQ ID NO:3 or the 4.5 *Pst*I/*Pst*I fragment of plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524 with genomic DNA isolated from an organism of interest, particularly from another monocot. The skilled artisan knows how to proceed to find such sequences as there are many ways known in the

art to identify homologous sequences from other organisms. Such newly identified DNA molecules then can be sequenced and the sequence can be compared to any one of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or to the nucleotide sequence of the 4.5 *Pst*I/*Pst*I fragment of pBSK+LOX4A which has been deposited under accession no DSM 13524, and tested for promoter activity. Within the scope of the present invention are DNA molecules having at least 75%, preferably 80%, more preferably 90%, and most preferably 95% sequence identity to the nucleotide sequence of any one of SEQ ID NOs:1, 2, or 3 over a length of at least 50 nucleotides. The percentage of sequence identity is determined using computer programs that are based on dynamic programming algorithms. Computer programs that are preferred within the scope of the present invention include the BLAST (Basic Local Alignment Search Tool) search programs designed to explore all of the available sequence databases regardless of whether the query is protein or DNA. Version BLAST 2.0 (Gapped BLAST) of this search tool has been made publicly available on the Internet (currently <http://www.ncbi.nlm.nih.gov/BLAST/>). It uses a heuristic algorithm which seeks local as opposed to global alignments and is therefore able to detect relationships among sequences which share only isolated regions. The scores assigned in a BLAST search have a well-defined statistical interpretation. Said programs are preferably run with optional parameters set to the default values.

If desired, the promoters of the present invention can be fused with the nucleotide sequence encoding a transit peptide according to the invention for example, by using the nucleotide sequence depicted in SEQ ID NO:4, for chemically regulated expression of an associated coding region of interest in plastids, particularly in chloroplasts.

A chemical regulator according to the invention is defined as a substance which regulates expression of a gene through a chemically regulatable DNA sequence. The substance, in ionic or neutral form, with or without solvating or other complexing molecules or anions, will usually be exogenous relative to the system containing the chemically regulatable gene at the time regulation is desired. The use of exogenous chemical regulators is preferred because of the ease and convenience of controlling the amount of regulator in the system. However, the invention also includes the use of endogenous regulators, e.g., chemicals whose activities or levels in the system are artificially controlled by other components in, or acting on, the system.

Chemical regulators according to the invention include benzoic acid, salicylic acid, polyacrylic acid and substituted derivatives thereof; suitable substituents include lower alkyl, lower alkoxy, lower alkylthio and halogen, but particularly INA, BTH, probenazole, jasmonate, and methyl jasmonate.

An additional group of regulators for the chemically regulatable DNA sequences and chimeric genes of this invention is based on the benzo-1,2,3-thiadiazole structure and includes, but is not limited to, the following types of compounds: benzo-1,2,3-thiadiazolecarboxylic acid, benzo-1,2,3-thiadiazolethiocarboxylic acid, cyanobenzo-1,2,3-thiadiazole, benzo-1,2,3-thiadiazolecarboxylic acid amide, benzo-1,2,3-thiadiazolecarboxylic acid hydrazide, and derivatives thereof.

A preferred group of regulators includes, but is not limited to, benzo-1,2,3-thiadiazole-7-carboxylic acid, benzo-1,2,3-thiadiazole-7-thiocarboxylic acid, 7-cyanobenzo-1,2,3-thiadiazole, benzo-1,2,3-thiadiazole-7-carboxylic acid amide, benzo-1,2,3-thiadiazole-7-carboxylic acid hydrazide, and derivatives thereof.

Suitable derivatives encompass but are not limited to representatives of said types of compounds wherein the benzo-1,2,3-thiadiazole moiety is unsubstituted or substituted by small substituents normally used in aromatic ring systems of agrochemicals such as lower alkyl, lower alkoxy, lower haloalkyl, lower haloalkoxy, lower alkylthio, cyano, nitro and halogen. Suitable derivatives further encompass, but are not limited to, representatives of said benzo-1,2,3-thiadiazole compounds wherein either the carboxylic acid, the thiocarboxylic acid, the carboxylic acid amide or the carboxylic acid hydrazide functional group is unsubstituted or substituted by aliphatic, araliphatic or aromatic residues. Suitable residues encompass, but are not limited to, alkyl (especially lower alkyl), alkoxy (especially lower alkoxy), lower alkoxyalkyl, alkoxyalkoxyalkyl, cycloalkyl, cycloalkylalkyl, phenylalkyl (especially benzyl), naphthylalkyl, phenoxyalkyl, alkenyl, and alkynyl, wherein the alkyl part of the substituent is unsubstituted or substituted by hydroxy, halogen, cyano or nitro, and the aromatic part of the substituent is unsubstituted or substituted by small substituents normally used in aromatic ring systems in agrochemicals such as lower alkyl, lower alkoxy, lower haloalkyl, lower haloalkoxy, lower alkylthio, cyano, nitro and halogen.

Regulators based on the benzo-1,2,3-thiadiazole structure encompass all molecular systems capable of releasing the molecule actually acting as the regulator.

A preferred group of regulators based on the benzo-1,2,3-thiadiazole structure includes benzo-1,2,3-thiadiazole-carboxylic acid, alkyl benzo-1,2,3-thiadiazolecarboxylate in which the alkyl group contains one to six carbon atoms, and substituted derivatives of these

compounds. Suitable substituents include lower alkyl, lower alkoxy, lower alkylthio and halogen. In particular, benzo-1,2,3-thiadiazole-7-carboxylic acid and its alkyl esters, e.g. methyl benzo-1,2,3-thiadiazole-7-carboxylate, are preferred inducers for the chimeric DNA sequences comprising chemically regulatable DNA sequences isolated from PR protein genes. The syntheses of the mentioned chemical regulators and their utility as biocides may be discerned from British Patent 1,176,799 and Kirby, P. et al., J. Chem. Soc. C 2250 (1970).

Among the preferred species based on the benzo-1,2,3-thiadiazole structure there may be mentioned, for example, benzo-1,2,3-thiadiazole-7-carboxylic acid, methyl benzo-1,2,3-thiadiazole-7-carboxylate, n-propyl benzo-1,2,3-thiadiazole-7-carboxylate, benzyl benzo-1,2,3-thiadiazole-7-carboxylate, benzo-1,2,3-thiadiazole-7-carboxylic acid sec-butylhydrazide, and the like.

An additional group of regulators for the chemically regulatable DNA sequences of this invention is based on the pyridine carboxylic acid structure, such as the isonicotinic acid structure and preferably the haloisonicotinic acid structure. Preferred are dichloroisonicotinic acids and derivatives thereof, for example the lower alkyl esters. Suitable regulators of this class of compounds are, for example, 2,6-dichloroisonicotinic acid, and the lower alkyl esters thereof, especially the methyl ester.

The chemical regulators may be applied in pure form, in solution or suspension, as powders or dusts, or in other conventional formulations used agriculturally or in bioreactor processes. Such formulations may include solid or liquid carriers, that is, materials with which the regulator is combined to facilitate application to the plant, tissue, cell or tissue culture, or the like, or to improve storage, handling or transport properties. Examples of suitable carriers include silicates, clays, carbon, sulfur, resins, alcohols, ketones, aromatic hydrocarbons, and the like. If formulated as a conventional wettable powder or aqueous emulsion, the regulator formulation may include one or more conventional surfactants, either ionic or non-ionic, such as wetting, emulsifying or dispersing agents.

The regulators may also be applied to plants in combination with another agent which is desired to afford some benefit to the plant, a benefit related or unrelated to the trait controlled by any chimeric gene which is regulated by the regulator. For example, a regulator can be admixed with a fertilizer and applied just before the expression of a transgenic trait unrelated to fertilization is desired. Or it can be combined with a herbicide and applied to mitigate the effect of the herbicide at the time when such effect would otherwise be at a maximum.

As a liquid formulation the regulator may be applied as a spray to plant leaves, stems or branches, to seeds before planting or to the soil or other growing medium supporting the plant. Regulators can also be used in bioreactor systems, regulation being achieved by a single addition of regulator formulation to the reaction medium or by gradual addition over a predetermined period of time.

The regulator is applied in an amount and over a time sufficient to effect the desired regulation. A preferred regulator is one which shows no, or only minimal phytotoxic or other deleterious effect on the plant, plant tissue or plant cells to which it is applied in the amount applied.

A further aspect of the invention is a process for regulating transcription of a chemically inducible, but not wound or pathogen inducible gene, which process comprises applying such a chemical regulator to plant tissue, plant or seed containing a chemically regulatable nucleotide sequence as described *supra*. Preferred is such a process wherein the plant tissue, plant or seed contains a chemically regulatable nucleotide sequence mentioned above as being preferred.

It is another object of the present invention to provide recombinant nucleic acid molecules comprising a promoter according to the invention operably linked to a nucleotide sequence of interest. The nucleotide sequence of interest can, for example, code for a ribosomal RNA, an antisense RNA or any other type of RNA that is not translated into protein. In another preferred embodiment of the invention the nucleotide sequence of interest is translated into a protein product. The nucleotide sequence associated with the promoter sequence may be of homologous or heterologous origin with respect to the plant to be transformed. The sequence may also be entirely or partially synthetic. Regardless of the origin, the associated nucleotide sequence will be expressed in the transformed plant in accordance with the expression properties of the promoter to which it is linked. In case of homologous nucleotide sequences associated with the promoter sequence, the promoter according to the invention is useful for ectopic expression of said homologous sequences. Ectopic expression means that the nucleotide sequence associated with the promoter sequence is expressed in tissues and organs and/or at times where said sequence may not be expressed under control of its own promoter. In one particular embodiment of the invention, expression of nucleotide sequence associated with the promoter sequence

suppresses its own expression and that of the original copy of the gene by a process called cosuppression.

In a preferred embodiment of the invention the associated nucleotide sequence may code for a protein that is desired to be expressed in a chemically inducible, but not wound- or pathogen inducible fashion. Such nucleotide sequences preferably encode proteins conferring a desirable phenotypic trait to the plant transformed therewith. Examples are nucleotide sequences encoding proteins conferring antibiotic resistance, virus resistance, insect resistance, disease resistance, or resistance to other pests, herbicide tolerance, improved nutritional value, improved performance in an industrial process or altered reproductive capability. The associated nucleotide sequence may also be one that is transferred to plants for the production of commercially valuable enzymes or metabolites in the plant. Embraced by the present invention are also selectable or screenable marker genes, i.e. genes comprising a nucleotide sequence of the invention operably linked to a coding region encoding a selectable or screenable trait.

Examples of selectable or screenable marker genes are described below. For certain target species, different antibiotic or herbicide selection markers may be preferred. Selection markers used routinely in transformation include the *npII* gene which confers resistance to kanamycin, paromomycin, geneticin and related antibiotics (Vieira and Messing, 1982, Gene 19: 259-268; Bevan et al., 1983, Nature 304:184-187) the bacterial *aadA* gene (Goldschmidt-Clermont, 1991, Nucl. Acids Res. 19: 4083-4089), encoding aminoglycoside 3'-adenylyltransferase and conferring resistance to streptomycin or spectinomycin, the *hph* gene which confers resistance to the antibiotic hygromycin (Blochliger and Diggelmann, 1984, Mol. Cell. Biol. 4: 2929-2931), and the *dhfr* gene, which confers resistance to methotrexate (Bourouis and Jarry, 1983, EMBO J. 2: 1099-1104). Other markers to be used include a phosphinothricin acetyltransferase gene, which confers resistance to the herbicide phosphinothricin (White et al., 1990, Nucl. Acids Res. 18: 1062; Spencer et al. 1990, Theor. Appl. Genet. 79: 625-631), a mutant EPSP synthase gene encoding glyphosate resistance (Hinchee et al., 1988, Bio/Technology 6: 915-922), a mutant acetolactate synthase (ALS) gene which confers imidazolinone or sulfonyleurea resistance (Lee et al., 1988, EMBO J. 7: 1241-1248), a mutant *psbA* gene conferring resistance to atrazine (Smeda et al., 1993, Plant Physiol. 103: 911-917), or a mutant protoporphyrinogen oxidase gene as described in EP 0 769 059. Selection markers resulting in positive selection, such as a phosphomannose isomerase gene, as described in patent application WO 93/05163, are also used.



Identification of transformed cells may also be accomplished through expression of screenable marker genes such as genes coding for chloramphenicol acetyl transferase (CAT),  $\beta$ -glucuronidase (GUS), luciferase (LUC), and green fluorescent protein (GFP) or any other protein that confers a phenotypically distinct trait to the transformed cell.

It is a further objective of the invention to provide recombinant expression vectors comprising a nucleotide sequence of the invention fused to an associated nucleotide sequence of interest. In these vectors, foreign nucleic acid molecules can be inserted into a polylinker region such that these exogenous sequences can be expressed in a suited host cell which may be, for example, of bacterial or plant origin. For example, the plasmid pBI101 derived from the *Agrobacterium tumefaciens* binary vector pBIN19 allows cloning and testing of promoters using beta-glucuronidase (GUS) expression signal (Jefferson et al, 1987, EMBO J 6: 3901-3907). The size of the vector is 12.2 kb. It has a low-copy RK2 origin of replication and confers kanamycine resistance in both bacteria and plants. There are numerous other expression vectors known to the person skilled in the art that can be used according to the invention.

It is a further objective of the present invention to provide transgenic plants comprising the recombinant DNA sequences of the invention. The invention thus relates to plant cells, to plants derived from such cells, to plant material, to the progeny and to seeds derived from such plants, and to agricultural products with improved properties obtained by any one of the transformation methods described below. Plants transformed in accordance with the present invention may be monocots or dicots and include, but are not limited to, rice, maize, wheat, barley, rye, sweet potato, sweet corn, bean, pea, chicory, lettuce, cabbage, cauliflower, broccoli, turnip, radish, spinach, asparagus, onion, garlic, pepper, celery, squash, pumpkin, hemp, zucchini, apple, pear, quince, melon, plum, cherry, peach, nectarine, apricot, strawberry, grape, raspberry, blackberry, pineapple, avocado, papaya, mango, banana, soybean, tomato, sorghum, sugarcane, sugar-beet, sunflower, rapeseed, clover, tobacco, carrot, cotton, alfalfa, potato, eggplant, cucumber, *Arabidopsis thaliana*, and woody plants such as coniferous and deciduous trees. Preferred plants to be transformed are rice, maize, wheat, barley, cabbage, cauliflower, pepper, squash, melon, soybean, tomato, sugar-beet, sunflower or cotton, but especially rice, maize, wheat, *Sorghum bicolor*, orchardgrass, sugar beet and soybean. The recombinant DNA sequences of the invention can be introduced into the plant cell by a number of well-known methods. Those skilled in the art will appreciate that the choice of such method might depend on the type of plant which is targeted for transformation, i.e., monocot or dicot. Suitable methods of

transforming plant cells include microinjection (Crossway et al., 1986, *Bio Techniques* 4:320-334), electroporation (Riggs and Bates, 1986, *Proc. Natl. Acad. Sci., USA* 83:5602-5606), *Agrobacterium*-mediated transformation (Hinchee et al., 1988, *Bio/Technology* 6:915-922; EP 0 853 675), direct gene transfer (Paszkowski et al., 1984, *EMBO J.* 3:2717-2722), and ballistic particle acceleration using devices available from Agracetus, Inc., Madison, Wisconsin and Dupont, Inc., Wilmington, Delaware (see, for example, US Patent No. 4,945,050 and McCabe et al., 1988, *Bio/Technology* 6:923-926). The cells to be transformed may be differentiated leaf cells, embryogenic cells, or any other type of cell. In the direct transformation of protoplasts, the uptake of exogenous genetic material into a protoplast may be enhanced by the use of a chemical agent or an electric field. The exogenous material may then be integrated into the nuclear genome. The previous work is conducted in dicot tobacco plants, which resulted in the foreign DNA being incorporated and transferred to progeny plants (Paszkowski et al., 1984, *EMBO J.* 3:2712-2722; Potrykus et al., 1985, *Mol. Gen. Genet* 199:169-177). Monocot protoplasts, for example, of *Triticum monococcum*, *Lolium multiflorum* (Italian rye grass), maize, and Black Mexican sweet corn, are transformed by this procedure. An additional preferred embodiment is the protoplast transformation method for maize as disclosed in EP 0 292 435, as well as in EP 0 846 771. For maize transformation also see Koziel et al., 1993, *Bio/Technology* 11:194-200. Transformation of rice can be carried out by direct gene transfer techniques utilizing protoplasts or particle bombardment. Protoplast-mediated transformation is described for Japonica-types and Indica-types (Zhang et al., 1988, *Plant Cell Rep.*, 7:379-384; Shimamoto et al., 1989, *Nature* 338:274-276; Datta et al., 1990, *Bio/Technology* 8:736-740). Both types described above are also routinely transformable using particle bombardment (Christou et al., 1991, *Bio/Technology* 9:957-962). Patent application No. EP 0 332 581 describes techniques for the generation, transformation and regeneration of *Pooideae* protoplasts. These techniques allow the transformation of all *Pooideae* plants including *Dactylis* and wheat. Furthermore, wheat transformation is described in patent application No. EP 0 674 715; and by Weeks et al., 1993 (*Plant Physiol.* 102:1077-1084). The thus-constructed plant expression vector can, for example, be introduced into the calli of rice according to the conventional plant transformation method, and the differentiation of roots and leaves is induced therefrom, and then, can be transferred to a flowerpot for cultivation, thereby obtaining the transformed rice.

The plants resulting from transformation with the DNA sequences or vectors of the present invention will express a nucleotide sequence of interest throughout the plant and in most tissues and organs.

The genetic properties engineered into the transgenic plants described above are passed on by sexual reproduction or vegetative growth and can thus be maintained and propagated in progeny plants. Generally said maintenance and propagation make use of known agricultural methods developed to fit specific purposes such as tilling, sowing or harvesting. Specialized processes such as hydroponics or greenhouse technologies can also be applied. Use of the advantageous genetic properties of the transgenic plants according to the invention can further be made in plant breeding that aims at the development of plants with improved properties such as tolerance of pests, herbicides, or stress, improved nutritional value, increased yield, or improved structure causing less loss from lodging or shattering. The various breeding steps are characterized by well-defined human intervention such as selecting the lines to be crossed, directing pollination of the parental lines, or selecting appropriate progeny plants. Depending on the desired properties different breeding measures are taken. The relevant techniques are well known in the art and include but are not limited to hybridization, inbreeding, backcross breeding, multiline breeding, variety blend, interspecific hybridization, aneuploid techniques, etc. Hybridization techniques also include the sterilization of plants to yield male or female sterile plants by mechanical, chemical or biochemical means. Cross pollination of a male sterile plant with pollen of a different line assures that the genome of the male sterile but female fertile plant will uniformly obtain properties of both parental lines. Thus, the transgenic plants according to the invention can be used for the breeding of improved plant lines that for example increase the effectiveness of conventional methods such as herbicide or pesticide treatment or allow to dispense with said methods due to their modified genetic properties. Alternatively new crops with improved stress tolerance can be obtained that, due to their optimized genetic "equipment", yield harvested product of better quality than products that were not able to tolerate comparable adverse developmental conditions.

It is another objective of the present invention to provide nucleotide sequences that can be used to express a nucleotide sequence of interest in a desired organism. Such molecules are commonly referred to as "promoters." This organism can be a bacterium, a plant or any other organism of interest.

Furthermore, the disclosure of SEQ ID NOs:1 to 3 enables a person skilled in the art to design oligonucleotides for polymerase chain reactions which attempt to amplify DNA

fragments from templates comprising a sequence of nucleotides characterized by any continuous sequence of 15 and preferably 20 to 30 or more base pairs in SEQ ID NOs:1, 2, or 3. Said nucleotides comprise a sequence of nucleotides which represents 15 and preferably 20 to 30 or more base pairs of SEQ ID NOs:1, 2, or 3. Polymerase chain reactions performed using at least one such oligonucleotide and their amplification products constitute another embodiment of the present invention.

#### **BRIEF DESCRIPTION OF THE SEQUENCES IN THE SEQUENCE LISTING**

SEQ ID NO:1	part of the 5' upstream sequence of the rice RCI-1 gene
SEQ ID NO:2	part of the rice RCI-1 gene including putative TATA box and putative start codon
SEQ ID NO:3	part of the rice RCI-1 gene including part of intron 1, exon 2, intron 2 and part of exon 3
SEQ ID NO:4	nucleotide sequence of the rice lipoxygenase RCI-1 transit peptide
SEQ ID NO:5	nucleotide sequence of the rice lipoxygenase RCI-1 cDNA
SEQ ID NO:6	amino acid sequence of the rice lipoxygenase RCI-1 transit peptide
SEQ ID NO:7	deduced amino acid sequence of the rice lipoxygenase RCI-1 cDNA
SEQ ID NO:8	degenerate primer
SEQ ID NO:9	anchored oligo dT reverse primer
SEQ ID NO:10	forward primer
SEQ ID NO:11	reverse primer
SEQ ID NO:12	reverse primer R1
SEQ ID NO:13	forward primer F1
SEQ ID NO:14	forward primer F2
SEQ ID NO:15	primer attB1
SEQ ID NO:16	primer attB2
SEQ ID NO:17	nucleotide sequence of the about 4.5 kb <i>Pst</i> I/ <i>Pst</i> I fragment from plasmid pBSK+LOX4a
SEQ ID NO:18	part of the 5' upstream sequence of the rice RCI-1 gene obtained by PCR with forward primer F1 and reverse primer R1

SEQ ID NO:19	part of the 5' upstream sequence of the rice RCI-1 gene obtained by PCR with forward primer F2 and reverse primer R1
SEQ ID NO:20	Gateway™ modified pNOV2347 binary Gateway™ destination vector with GIG reporter gene
SEQ ID NO:21	GIG, GUS intron GUS, GUS coding sequence with intron
SEQ ID NO:22	pNOV6800 binary vector

Deposit

Deposited material	Accession number	Date of deposit
pBSK+LOX4A	DSM 13524	06.06.2000
pNOV6800	NRRL B-30480	May 25, 2001

The deposit of pBSK + LOX4A was made with the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Mascheroder Weg 1b, D-38124 Braunschweig, Deutschland. The deposit of pNOV6800 was made with the Agricultural Research Service Culture Collection (NRRL), of the National Center for Agricultural Utilization Research, Agricultural Research Service, United States Department of Agriculture, 1815 North University Street, Peoria, Illinois 61604 USA.

Below are illustrative examples of the present invention. The present invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and any constructs, promoters, transit peptides or enzymes which are functionally equivalent are within the scope of this invention.

**EXAMPLES**

Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described, for example, by Sambrook et al., 1989, "Molecular Cloning", Cold Spring Harbor, Cold Spring Harbor Laboratory Press, NY and by Ausubel et al., 1994, "Current protocols in molecular biology", John Wiley and Sons, New York.

**Example 1: Plant material and treatment**

Rice plants (*Oryza sativa* cv. Kusbue) are grown in pots with clay soil that are soaked with an iron fertilizer solution (Gesal Pflanzen Tonic, Novartis, Basel, Switzerland) under a 16 h light/8 h dark cycle at 25 °C and 80% humidity.

*M. grisea* (race 007 and 031 from the Institute of Biochemistry, Faculty of Agriculture, Tamagawa University, Machida-shi, Tokyo 194, Japan) is cultivated on oat-meal starch agar (30 g l<sup>-1</sup> oat-flakes, 20 g l<sup>-1</sup> agar-agar, 10 g l<sup>-1</sup> starch and 2 g l<sup>-1</sup> yeast extract). After incubation at 27 °C for 2 weeks, aerial mycelia are removed with a sterile spatula and synchronous sporulation is induced by further incubation under black light (310-360 nm). For inoculations the concentration of conidia is adjusted to 1 x 10<sup>6</sup> ml<sup>-1</sup> in a spraying solution (1 g l<sup>-1</sup> gelatine, 0.1 % Tween-20).

Plants are inoculated 12 - 14 days after sowing by spraying the conidial suspension onto the leaves. After an incubation for 24 h in a dark moist chamber (26 °C, approx. 100 % relative humidity), plants are kept in a humid atmosphere under the same temperature and light regime as described above.

Plant treatment with the chemical inducers is done 10 days after sowing at the emergence of leaf 3. All chemical concentrations are given as ppm (mg active ingredient l<sup>-1</sup> of applied solution). Probenazole is applied as a 250 ppm solution of the pure substance by soil drench as described (Thierson et al. (1995) Systemic acquired resistance in rice: Studies on the mode of action of diverse substances inducing resistance in rice to *Pyricularia oryzae*.

*Mededelingen Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen Universiteit Gent*. 60, 421-430). Formulations of BTH (1:1 (w/w) mixture of active ingredient and wettable powder) and INA (1:3 (w/w) mixture of active ingredient and wettable granulate) are applied onto leaves by spraying. All controls are done by application of spray-solutions without active substance. Jasmonic acid is applied as a 1 mM solution in ethanol as described (Schweizer et al. (1997) *Plant Physiol.* 114, 79-88). Wounding and measurement of gene expression in systemic tissue is done according to (Schweizer et al. (1998) *Plant J.* 14, 475-481).

#### **Example 2: cDNA library construction**

Total RNA is extracted from rice leaves treated with 100 ppm INA and harvested after 24 and 48 hours. PolyA<sup>+</sup>-RNA is prepared as described in Example 2 and equal amounts from both time points are pooled. A cDNA library is constructed using the lambda Zap Express cDNA Synthesis Kit (Stratagene, La Jolla, CA) according to the manufacturer's instructions.

**Example 3: Cloning of the rice lipoxygenase cDNA RCI-1****A. Cloning of a rice lipoxygenase cDNA fragment by PCR**

A PCR-based strategy is used to generate a lipoxygenase cDNA fragment from INA-treated rice leaves. By aligning two lipoxygenase sequences from rice (Peng et al. (1994) *J. Biol. Chem.* 269, 3755-3761; Ohta et al. (1992) *Eur. J. Biochem.* 206, 331-336) and one from wheat (Görlach et al. (1996) *Plant Cell.* 8, 629-643), several conserved regions are identified, one of which is near the C-terminus and contains the amino acid sequence HAAVNFG that is invariant in all three sequences.

Total RNA is extracted as described (Dudler & Hertig (1992) *J. Biol. Chem.* 267, 5882-5888) from untreated control leaves and from leaves sprayed with a 100 ppm INA solution 24 and 48 hours after treatment. Poly A<sup>+</sup>-RNA is prepared from total RNA using the quick mRNA isolation kit from Stratagene (La Jolla, CA). The polyA<sup>+</sup>-mRNA samples from the two time points are pooled and 1 µg aliquots of poly A<sup>+</sup> RNA are used as templates for RT-PCR using the degenerate oligonucleotide 5'-CAYGCNGTNAANTTYGG-3' (SEQ ID NO:8), which corresponds to the HAAVNFG amino acid sequence motif in the C-terminal region of the rice RLL2 lipoxygenase (Peng et al. (1994) *J. Biol. Chem.* 269, 3755-3761) as the forward primer and an anchored oligo-dT reverse primer (5'-AATGCTTTTTTTTTTTTTT-3', SEQ ID NO:9).

Reverse transcription is done as follows:

1 µl RNA (1µg/ml)

8 µl 5x RT-buffer

4 µl anchored oligo-dT reverse primer (100 pmol/µl)

4 µl dNTP (10 mM each)

18 µl H<sub>2</sub>O<sub>DEPC</sub>

are mixed and incubated at 94°C for 15 min. This is followed by 1h incubation at 42°C. Five minutes after the transfer to 42°C, 3 µl AMV-RT (Boehringer) and 2 µl RNase Inhibitor (Boehringer) are added. Reverse transcription is terminated by 5 min incubation at 75°C.

PCR with reverse transcribed RNA is done as follows:

3 µl cDNA (see above)

10 µl 10x PCR-buffer

1 µl anchored oligo-dT reverse primer (100 pmol/µl)

1 µl degenerate primer (100 pmol/µl)

2 µl dNTP (10 mM each)

0,5 µl Taq DNA polymerase (Boehringer)

82,5 µl H<sub>2</sub>O

are mixed and subjected to the following program:

1 cycle    94°C/15 min    39°C/ 2 min.    72°C/ 3 min

35 cycle   94°C/30 sec.    39°C/ 30 sec.    72°C/ 1 min

1 cycle    94°C/1 min 39°C/ 2 min.    72°C/ 10 min

Then, 0,5 µl fresh Taq DNA polymerase are added, and 20 more cycles are performed under the conditions given above.

PCR products derived from treated and untreated leaves are visualized on ethidiumbromide-stained agarose gels. A PCR product of approximately 600 bp arises only in the INA-treated sample but not in the control. The piece of gel corresponding to the about 600 bp band present only in the lane with INA-treated probes is cut out. The DNA is subsequently eluted from the gel and cloned into the pGEMT<sub>easy</sub> vector (Promega, Madison, USA) and the resultant plasmid designated pKL-5.

#### B. cDNA library screening to obtain a full-length rice lipoxygenase cDNA clone

The <sup>32</sup>P-labelled insert of pKL-5 is used as a probe to screen a lambda cDNA library constructed from INA-treated rice leaves (see Example 2). Positive clones are purified. The one with the largest insert is designated RCI-1 (rice chemically induced cDNA 1) and subcloned into the pBK-CMV (Stratagene) phagemid vector by in vivo excision according to the instructions of the manufacturer. The resulting plasmid is called pRCI-1. The RCI-1 insert is sequenced on both strands by primer walking using CY5-labelled primers and an ALF DNA-sequencer (Pharmacia, Uppsala, Sweden).

The *RCI-1* cDNA insert (SEQ ID NO:5) consists of 3018 bp and contains an open reading frame of 2766 bp (from base 48 to base 2816 of SEQ ID NO:5) encoding a protein of 922 amino acid residues (SEQ ID NO:7) with a predicted Mr of 105 kDa. The presumed translation initiation site is the first methionine codon in the open reading frame. Sequence comparison revealed that the RCI-1 protein was most similar to the barley LOX2:Hv:1 (Vörös et al. (1998) *Eur. J. Biochem.* 251, 36-44), showing 60% identity and 68% similarity at the amino acid level. Sequence similarity (identity) to the two already published rice lipoxygenase forms at the amino acid level were 52% (43%) in comparison to L-2 (Ohta et



al. (1992) *Eur. J. Biochem.* 206, 331-336) and 58% (50%) to RLL2 (Peng et al. (1994) *J. Biol. Chem.* 269, 3755-3761), respectively.

The RCI-1 rice lipoxygenase has an N-terminal extension (SEQ ID NO:6, corresponding to amino acid 1 to 36 of SEQ ID NO:7) that is thought to direct this class of proteins to plastids, in particular to chloroplasts. This putative chloroplast targeting sequence clearly separates this class of lipoxygenase (LOX) species from another class that is predominately found in kernels and seedlings and that includes LoxA, LoxB, and LoxC from barley (van Mechelen et al. (1999) *Plant Mol. Biol.* 39, 1283-1298), and LOX L-2 from rice (Ohta et al. (1992) *Eur. J. Biochem.* 206, 331-336).

#### **Example 4: Southern blot analysis**

Genomic DNA is extracted from rice leaves using a CTAB procedure (Ausubel et al. (1987) *Current protocols in molecular biology*, Wiley and Sons, New York). Digestion with restriction enzymes, electrophoretic separation on agarose gels, and transfer to GeneScreen membranes (Dupont NEN, Brussels, Belgium) are performed according to standard procedures. Filters are hybridized to a <sup>32</sup>P-labeled probe consisting of an *EcoRI/HindIII* fragment of pRCI-1 that contains the first 1280 bp of the *RCI-1* cDNA in 1 M NaCl, 1% SDS, 10% dextrane sulphate, and 100 µg ml<sup>-1</sup> denatured salmon sperm DNA overnight at 68° C. Filters are washed in 0.2 x SSC (1 x SSC is 150 mM NaCl; 15 mM sodium citrate); 0.1 % SDS at 65° C.

When DNA digested with a number of enzymes that do not cut within the sequence of the probe is analyzed, one to three bands hybridizing to the probe are detected. This suggests that in addition to *RCI-1*, there is at least one other rice gene that weakly crosshybridizes with the probe.

#### **Example 5: RCI-1 expression studies**

The effect of various stimuli on the abundance of *RCI-1* transcripts is investigated using RNA gel blot analysis. For this, total RNA is extracted from treated and untreated leaves as described (Dudler. & Hertig. (1992) *J. Biol. Chem.* 267, 5882-5888). For gel blot analysis, 10 µg of total RNA is loaded per slot and separated on formaldehyde agarose gels, transferred to GeneScreen membranes, and cross-linked using an UV crosslinker (Amersham, UK). Loading of the lanes is monitored by ethidium bromide staining of the gel before transfer. Filters are hybridized to a <sup>32</sup>P-labeled probe consisting of an *EcoRI/HindIII* fragment of

pRCI-1 that contains the first 1280 bp of the *RCI-1* cDNA in 1 M NaCl, 1% SDS, 10% dextrane sulphate, and 100 µg ml<sup>-1</sup> denatured salmon sperm DNA overnight at 68° C. Filters are washed in 0.2 x SSC (1 x SSC is 150 mM NaCl; 15 mM sodium citrate); 0.1 % SDS at 65° C. A 528 bp cDNA fragment encoding part of the rice ribosomal Protein L3 (RP-L3, accession number D12630) is used as a probe for a constitutively expressed transcript. This fragment is fortuitously amplified and cloned together with the partial lipoxygenase clone pKL-5. A time course experiment with rice leaves that have been treated with INA is analyzed by RNA gel blot analysis. The hybridization signal appears as a distinct band corresponding to an RNA of approximately 3200 bp length and a smeared signal of about 1200 to 1700 bp. By stripping and reprobing the same membrane with a constitutively expressed gene (RP-L3), the high quality of the RNA preparation is confirmed. The smeared signal thus indicates that *RCI-1* transcripts are particularly unstable, perhaps due to a high turn-over rate. The time course experiment reveals that *RCI-1* transcripts starts to accumulate 8 hours after treatment and reaches maximal levels after 24 to 48 h. Similarly, treatment with the resistance activator BTH, a functional analogue of INA, also induces *RCI-1* transcript accumulation as does the application of probenazole (tradename oryzemate), which represents a different class of resistance-inducing chemicals (Kessmann et al (1994) *Ann. Rev. Phytopathol.* 32, 439-459). The delayed time course of *RCI-1* mRNA accumulation in response to probenazole treatment may rather reflect the different mode of application, i. e. spraying onto the leaves in case of INA and BTH vs. soil drenching with probenazole, respectively, than a difference in signaling. Thus, *RCI-1* transcripts accumulate upon application of a number of different chemical resistance inducers. In contrast, *RCI-1* mRNA levels neither increase after inoculation with the non-host pathogen *P. syringae* pv. *syringae*, a biological inducer of resistance against rice blast (Smith & Métraux (1991) *Physiol. Mol. Plant Pathol.* 39, 451-461), nor upon infection with *M. grisea*, the causal agent of rice blast.

Furthermore, *RCI-1* transcript levels strongly increase 7 to 12 h after spraying of a 1 mM jasmonic acid (JA) solution onto rice leaves. Interestingly, wounding, which is known to increase endogenous levels of JA in rice and induces increased systemic protection against blast infection (Schweizer et al. (1998) *Plant J.* 14, 475-481; Schweizer et al. (1997) *Plant Physiol.* 114, 79-88), does not activate *RCI-1* transcription, neither locally nor systemically. To investigate whether the observed accumulation of *RCI-1* transcripts after treatment with chemical inducers correlates with the increase in lipoxygenase (LOX) enzyme activity in rice

leaves. To this end LOX activity is measured in the BTH-treated rice leaves that are also used for RNA gel blot analyses shown (see above). Consistent with the results of the RNA gel blot analysis, a significant increase in enzymatic activity is observed between 24 and 48 h after BTH treatment. In addition, the BTH dose that is sufficient to trigger *RCI-1* transcript accumulation also causes an increase in LOX enzyme activity. Both results are compatible with the assumption that the increase in enzyme activity is predominantly due to the activation of the *RCI-1* (and homologous) gene(s). To analyze this hypothesis further, RNA derived from BTH-treated rice plants is probed with other rice LOX cDNAs that correspond to a pathogen-induced gene (*RLL2*; Peng et al. (1994) *J. Biol. Chem.* 269, 3755-3761) and a gene expressed in seedlings (*L-2*; Ohta et al. (1992) *Eur. J. Biochem.* 206, 331-336). In contrast to *RCI-1* mRNA, *RLL2* and *L-2* transcripts do not accumulate after treatment with INA, BTH, and probenazole.

#### **Example 6: Expression of RCI-1 in *E. coli***

The *RCI-1* coding region is placed under the control of an IPTG-inducible promoter of an *E. coli* expression vector. More specifically, the *RCI-1* cDNA is cloned into the pDS56/RBSII, *SphI* expression vector (Stüber et al. (1990) System for high-level production in *Escherichia coli* and rapid purification of recombinant proteins: Applications to epitope mapping, preparation of antibodies, and structure-function analysis. In *Immunological Methods* (Levkovits, I. & Pernis, B., eds) pp. 121-152. Academic Press, New York) from which the unique *PstI* site is eliminated by *PstI* digestion and religation after blunting the sticky ends using T4 DNA polymerase. The new vector is named pDS56/RBSII, *SphI* (–*PstI*). An *SphI* site is introduced at the translation initiation site of the *RCI-1* cDNA by PCR amplification of a 146 bp *RCI-1* fragment using the forward primer 5'-GTCAGCATGCTCACGGCCAC-3' (SEQ ID NO:10; the *SphI* site is underlined; the translation initiation codon is given in italics) and the reverse primer 5'- CATTGACGACCTCCGACAAG-3' (SEQ ID NO:11), which anneals downstream of an internal *XhoI* site (nucleotide position 149 of SEQ ID NO:5). The amplified fragment is cut with *SphI* and *XhoI* and ligated together with the 2.3 kb *XhoI/BamHI* fragment containing the middle part of the *RCI-1* cDNA (nucleotide position 149 to 2468 of (SEQ ID NO:5) in a single reaction into pDS56/RBSII, *SphI* (–*PstI*) that has been digested with *SphI* and *BamHI*. The resulting vector (pExpr1) is checked by restriction analysis. pExpr1 is then cut with *PstI* (at position 891 in the top strand of the cDNA insert, corresponding to base 891 of SEQ ID NO:5) and *SalI* (in the multiple cloning site of

pDS56/RBSII, *SphI* (–*PstI*) downstream of the insert), and the resulting cDNA fragment is replaced with the corresponding *PstI/XhoI* fragment of pRCI-1 (*XhoI* cleaves in the multiple cloning site downstream of the cDNA insert). The resulting construct (pExprRCI-1), which contains the complete RCI-1 coding region under the control of an IPTG-inducible promoter, is subsequently transformed into M15 *E. coli* cells (Stüber et al. (1990) System for high-level production in *Escherichia coli* and rapid purification of recombinant proteins: Applications to epitope mapping, preparation of antibodies, and structure-function analysis. In *Immunological Methods* (Levkovits, I. & Pernis, B., eds) pp. 121-152. Academic Press, New York).

Production of recombinant RCI-1 protein is induced by addition of 1 mM IPTG (final concentration) to a 250 ml *E. coli* M15 culture grown to an OD<sub>600</sub> of 1 and further incubation on a shaker overnight at 19°C. The cells are harvested by centrifugation (5000 g, 10 minutes) and resuspended in 5 ml lysis buffer (50 mM Na-phosphate buffer pH 7.5 containing 1 mg l-1 lysozyme). After a 30 minute incubation on ice, the lysate is centrifuged (12000 g, 15 minutes) and the pellet is transferred into a mortar and ground in extraction buffer (0.1 M K-phosphate buffer pH 7, 30 mg polyvinyl-poly-pyrrolidone, 1 mM EDTA). After centrifugation, the clear supernatant is used as an enzyme preparation for further biochemical analysis.

SDS-PAGE analysis of extracts of *E. coli* transformed with this construct reveals a novel protein with a molecular mass of about 103 kDa which is recognized by a LOX specific antibody on a western-blot. This size is compatible with the predicted value of 105 kDa (see Example 3).

#### **Example 7: Biochemical analysis of RCI-1 lipxygenase activity**

Lipxygenase activity is measured at 30°C photospectrometrically at 234 nm using linoleic acid as a substrate and 5-20 µl from the recombinant RCI-1 enzyme extract (Bohland et al. (1997) *Plant Physiol.* 114, 679-685). For determination of the pH optimum, different buffers with overlapping pH ranges (pH 4-6: 0.1 M Na-acetate; pH 6-8: 0.1 M Na-phosphate; pH 8-10: 0.1 M Tris-HCl) are used. The molar extinction coefficient of the reaction product,  $2.5 \times 10^7 \text{ cm}^{-1} \text{ mol}^{-1}$ , is used for the calculation of the enzyme activity.

Crude extracts of these bacteria exhibit increased LOX activity using linoleic acid as a substrate, while control extracts of *E. coli* without expression construct or containing the empty vector do not have detectable LOX activity. Maximal activity is observed around pH 8

to 9, indicating that RCI-1 must be classified as a type 1 LOX (Siedow (1991) *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 42, 145-188). However, it should be noted that recently a second classification based on the presence of a plastomic transit peptide is introduced (Shibata et al. (1994) *Plant Mol. Biol. Rep.* 12, 41-42). According to this scheme, RCI-1 must be classified as a type 2 LOX.

The products of the enzymatic activity of the RCI-1 protein are analyzed by HPLC (Bohland et al. (1997) *Plant Physiol.* 114, 679-685). Approximately 0.2 nkat enzyme activity obtained from recombinant RCI-1 protein is incubated with 50  $\mu$ l substrate solution (10  $\mu$ l of linoleic or  $\alpha$ -linolenic acid, respectively; 20  $\mu$ l ethanol, 20  $\mu$ l H<sub>2</sub>O) in 2 ml 0.1 M Tris-HCl pH 8.8 for 20 min. at 30°C. The reaction is stopped by lowering the pH to 3.0 with diluted HCl, and the hydroperoxides are extracted with 1 ml of CHCl<sub>3</sub> followed by two washes with water. The reaction products are subjected to HPLC-analysis (4- $\mu$ m particle size, Suprasphere-Si, 4.6 x 125 mm; Merck, Darmstadt, Germany). Isocratic elution is performed with hexane:2-propanol:acetonitrile:acetic acid (98.3:1.5:0.1:0.1, v/v/v/v) at a flow rate of 1 ml min<sup>-1</sup>. Products are detected at 234 nm and standards are obtained from Biomol (Hamburg, Germany) or prepared from linoleic or  $\alpha$ -linolenic acid by incubation with soybean lipoxygenase, respectively, as described (Bohland et al. (1997) *Plant Physiol.* 114, 679-685).

When the reaction products of recombinant RCI-1 are analyzed by HPLC, (13S)-hydroperoxy-(9Z, 11E, 15Z)-octadecatrienoic acid (13-HPOD) is the predominant product, irrespective of whether linoleic acid or linolenic acid served as a substrate for the enzyme. (9S)-hydroperoxy-(10E, 12Z, 15Z) octadecatrienoic acid (9-HPOD) is only detected in minor amounts.

#### **Example 8: RCI-1 transit peptide::GFP reporter-gene construction and transformation**

A chimeric gene is constructed that encodes a fusion protein containing the N-terminal 37 amino acids of the RCI-1 protein (SEQ ID NO:6) followed by four amino acids resulting from the cloning procedure followed by GFP sequence. To this end, pRCI-1 (see Example 3 B) is digested with *Xho*I, which cuts the top strand after position 149 of the cDNA insert (corresponds to base 149 of SEQ ID NO:5) and in the multiple cloning site downstream of the insert, and religated. The resulting plasmid contains the first 158 bp of the *RCI-1* cDNA, since the nucleotide sequence of the vector downstream of the cloning site is identical to base 150 to base 158 of SEQ ID NO:5. This plasmid is referred to as pRCI158. Its insert

comprising the transit peptide cDNA (SEQ ID NO:4) is cleaved out with *EcoRI* and *XbaI*, which both cut in the multiple cloning site, and the sticky ends are blunted by filling them in with Klenow enzyme. The blunted fragment is cloned into the filled-in and dephosphorylated *SpeI* cloning site of the binary pCambia 1302 vector (Cambia, Canberra, Australia), which contains the mGFP-5 coding region (MRC Laboratory of Molecular Biology, Cambridge, England). The correct orientation of the inserted fragment is checked by restriction digestion and the final construct is verified by sequencing. The construct is designated pRCI transit peptide::GFP and transformed into the *Agrobacterium tumefaciens* strain LBA 4404 by triparental mating. Transformation of *Arabidopsis* leaf cells is achieved by infiltration of *Agrobacterium* into intact leaves of *Arabidopsis thaliana*, ecotype Wassiljewskija (Ws), 14 days after germination according to Kapila et al. (1997) (*Plant Sci.* 122, 101-108).. Expression of the fusion protein is monitored 2 days after transformation by confocal imaging using a Leica-TCS confocal laser scanning microscope and a PLAPO x100 oil immersion objective (Leica Microsystems, Heidelberg, Germany) with the following filter settings: excitation 476/488 nm; GFP-emission 515-552 nm, chlorophyll-emission 673-695 nm. GFP fluorescence and chlorophyll autofluorescence are recorded simultaneously using independent 2-channel-detection.

Confocal imaging of leaves from transgenic *Arabidopsis* plants expressing the pRCI transit peptide::GFP construct reveals a strict congruence of GFP fluorescence and chlorophyll autofluorescence, indicating that the fusion protein is localized in the chloroplasts.

#### **Example 9: Cloning of the RCI-1 promoter region**

##### **A. Screening of a $\lambda$ -DASH genomic DNA library of rice**

A  $\lambda$ -DASH II/*Bam*HI DNA library representing genomic DNA derived from *Oryza sativa* cv. Norin plants is constructed according to the protocol of Stratagene (La Jolla, USA). The titer of the library is determined to be  $2.12 \times 10^{10}$  pfu/ml. Screening of the library is carried out following the protocol of Stratagene. The library is plated on four 530 cm<sup>2</sup> bio-assay dishes (Nalge Nunc Int., Naperville, USA) containing NZY agar. The density is adjusted to 150'000 pfu/plate and plating is carried out with *E. coli* XL1-blue MRA (Stratagene, La Jolla, USA) as a host strain according to the protocol of Stratagene. The plaques are transferred onto a nylon membrane (Hybond<sup>TM</sup>N 0.45  $\mu$ m, Amersham, Uppsala, Sweden) and the DNA is crosslinked in a UV crosslinker (Amersham, Uppsala, Sweden). A 900 bp *Pst*I-fragment representing the 5'-prime end of the rice *RCI-1* lipoygenase cDNA clone pRCI-1 (SEQ ID

NO:5) is labeled with  $^{32}\text{P}$  and hybridized overnight at 65°C to the plaque lifts according to standard procedures (Maniatis *et al.*, 1982). Two additional rounds of screening resulted in a positive  $\lambda$ -clone ( $\lambda\text{LOX4}$ ).

#### B. Subcloning of the putative LOX promoter region

Liquid lysate DNA preparations of the  $\lambda$ -clone are prepared according to standard procedures and analyzed by digestion with the *Pst*I restriction enzyme and gel electrophoresis on a 0.6 % (w/v) agarose gel. Southern blotting and subsequent hybridization of the membrane to the 900 bp *Pst*I-fragment of the *RCI-1* cDNA are done according to standard procedures. A strong band corresponding to a 4.5 kb fragment of  $\lambda\text{LOX4}$  is detected. The 4.5 kb DNA fragment is subcloned into a pBluescript/SK+ vector (Stratagene, La Jolla, USA). Transformation of *E. coli* strain DH5 $\alpha$  cells is done according to standard procedures and transformants are selected on LB Agar containing Ampicillin (100  $\mu\text{g/ml}$ ). This resulting clone is designated pBSK+LOX4A. Clone pBSK+LOX4A is deposited with the DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH on June 6, 2000 with accession number DSM 13524. Clone pBSK+LOX4A contains the *RCI-1* lipooxygenase promoter on a 4.5 kb *Pst*I/*Pst*I fragment and is further analyzed by DNA sequencing. Clone pBSK+LOX4A comprises, in a 5' to 3' direction, the nucleotide sequences depicted in SEQ ID NOs:1, 2 and 3. SEQ ID NO:1 comprises the 5'-end of the 4.5 kb *Pst*I/*Pst*I fragment. This nucleotide sequence is 358 nucleotides in length and contains at its 5' end in position 1 to 6 the *Pst*I-site. The region between SEQ ID NO:1 and SEQ ID NO:2 of the 4.5 kb *Pst*I/*Pst*I fragment is between about 240 and 440 bp in length. The central region of the 4.5 kb *Pst*I/*Pst*I fragment is shown in SEQ ID NO:2 and is 2104 bp in length. It contains the putative TATA box (position 1261 to 1266 SEQ ID NO:2), the putative start codon (position 1359 to 1361 of SEQ ID NO:2), as well as the 5' untranslated region and nucleotide sequences upstream of the putative TATA box. Comparison of the genomic DNA (SEQ ID NO:2) and the cDNA (SEQ ID NO:5) shows that the sequences located at position 1312 to 1701 of SEQ ID NO:2 comprise all or part of exon 1, and the sequences located at position 1702 to 2104 of SEQ ID NO:2 are the 5' part of intron 1. The region between SEQ ID NO:2 and SEQ ID NO:3 of the 4.5 kb *Pst*I/*Pst*I fragment is between about 85 and 285 bp in length. The 3' end of the 4.5 kb *Pst*I/*Pst*I fragment is shown in SEQ ID NO:3. This sequence depicts a nucleotide sequence of 1516 bp in length. It contains, in a 5' to 3' direction, the 3' end of intron 1 (position 1 to 97 of SEQ ID NO:3) followed by exon

2 (position 98 to 366 of SEQ ID NO:3), intron 2 (position 367 to 1283 of SEQ ID NO:3) and part of exon 3 (position 1284 to 1516 of SEQ ID NO:3). The *Pst*I site is located at position 1511 to 1516.

**Example 10: Plasmid amplification and DNA sequencing of the clone pBSK+LOX4A**

Transformants are grown at 37°C in a 50 ml over-night culture of LB Medium containing Ampicillin (100 µg/ml). Cells are harvested and plasmid DNA is extracted using the Jetstar Midi plasmid extraction kit (Genomed GmbH, Bad Oeynhausen, Germany).

Sequencing of the clone pBSK+LOX4A is carried out by the chain termination method (Maniatis *et al.* (1982) *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor). Sequencing reactions are performed the BigDye™ terminator cycle sequencing kit (Perkin-Elmer Corp., Norwalk, Connecticut) according to the instructions of the manufacturer and the sequences are determined with a 373 DNA-sequencer (Applied Biosystems, Foster City, California). They are assembled and analyzed using the Wisconsin Sequence Analysis package (Genetics Computer Group, Madison, Wisconsin). Ambiguities are clarified by comparison with the corresponding electropherogram print. SEQ ID NO:1 corresponds to the 5' end, SEQ ID NO:2 to the middle and SEQ ID NO:3 to the 3' end of the 4.5 kb insert of pBSK+LOX4A.

**Example 11: Preparation of a CaMV 35S promoter::RCI-1 cDNA construct**

The starting plasmid is the plant binary vector pBI 121 (Clontech, Palo Alto, California), which contains the *GUS* reporter gene under the control of the CaMV 35S promoter. The *GUS* reporter gene is removed from pBI 121 and replaced with the *RCI-1* cDNA. To this end, pBI 121 is digested with the restriction enzyme *Sst* I, and the sticky ends are filled in with dNTPs and T4 DNA polymerase according to standard procedures. After cutting with *Sma* I, the vector fragment is separated from the *GUS* reporter gene by agarose gel electrophoresis and religated. This vector is named pBI 121 (-GUS). pBI 121 (-GUS) is cut with *Bam* HI, the sticky ends are blunted by filling them in with dNTPs and T4 DNA polymerase, and the *RCI-1* cDNA fragment is ligated into this vector, after it has been cut out of pRCI-1 with *Eco* RI and *Xba* I and its sticky ends have been blunted with dNTPs and T4 DNA polymerase. After transformation into *E. coli*, plasmid is prepared from a number of colonies. The orientation of the *RCI-1* fragment is checked by restriction digestion using *Xba* I, which cuts immediately upstream of the insert, and *Bam* HI, which cuts the top strand



of the *RCI-1* cDNA after nucleotide 2688. The correct orientation results in a fragment of approx. 2700 bp in length, the wrong orientation in a fragment of approx. 350 bp. Then a plasmid which contains the *RCI-1* cDNA in the correct orientation, i. e. such that the filled in Eco RI site is next to the CaMV 35S promoter, is selected and designated p35Spromoter::*RCI-1*cDNA. For *Agrobacterium* mediated transformation, the plasmid is transformed into the *Agrobacterium tumefaciens* strain LBA 4404 by electroporation.

#### **Example 12: Transformation of rice**

The transformed *Agrobacterium tumefaciens* strain is grown for 3 days in the AB liquid medium supplemented with 30 mg/L hygromycin B and 3 mg/L tetracycline, and it is co-cultivated with three-week-old calli which are induced from the scutellum of mature seeds in the N6 medium (Chu, C.C. et al., Sci. Sin., 18, 659-668(1975)) containing 2 mg/L 2,5-D, on the 2N6-As medium supplemented with 1 mM betaine (Hiei, Y. et al., Plant J., 6, 271-282(1994)) in darkness at 25 °C for 2-3 days. The co-cultivated calli are washed with sterile water containing 100 mg/L cefotaxime, and again incubated on an N6 medium containing 40 mg/L hygromycin and 250 mg/L cefotaxime for 3 weeks. Actively growing hygromycin-resistant calli are transferred onto the selection medium [for example, MS media (Life Technologies) + 0.2 mg/L NAA (naphthalene acetic acid) + 2 mg/L kinetin + 2% sorbitol + 1.6% phytagar (Gibco) + 50 mg/L hygromycin B + 250 mg/L cefotaxime], and then cultivated for 2-3 weeks under continuous light condition of 40  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . The thus-obtained plantlets are potted and grown in a growth chamber under 10h light/14h dark condition to obtain transgenic rice plants.

#### **Example 13: Transformation of maize**

Type I embryogenic maize callus cultures (Green *et al*, Miami Winter Symposium 20,1983) are initiated from immature embryos, 1.5 - 2.5 mm in length, from greenhouse grown material. Embryos are aseptically excised from surface-sterilized ears approximately 14 days after pollination. Embryos may be placed on D callus initiation media with 2% sucrose and 5 mg/L chloramben (Duncan *et al*, Planta 165: 322-332,1985) or onto KM callus initiation media with 3% sucrose and 0.75 mg/L 2,4-d (Kao and Michayluk, Planta 126:105-110, 1975). Embryos and embryogenic cultures are subsequently cultured in the dark. Embryogenic responses are removed from the explants after ~14 days. Embryogenic responses from D callus initiation media are placed onto D callus maintenance media with

2% sucrose and 0.5 mg/L 2,4-d while those from KM callus initiation media are placed onto KM callus maintenance media with 2% sucrose and 5 mg/L Dicamba. After 3 to 8 weeks of weekly selective subculture to fresh maintenance media, high quality compact embryogenic cultures are established. Actively growing embryogenic callus pieces are selected as target tissue for gene delivery. The callus pieces are plated onto target plates containing maintenance medium with 12% sucrose approximately 4 hours prior to gene delivery. The callus pieces are arranged in circles, with radii of 8 and 10 mm from the center of the target plate.

Plasmid DNA containing the promoter-RCI-1-cDNA construct or a RCI-1-promoter-reporter gene construct is precipitated onto gold microcarriers as described in the DuPont Biolistics manual. Two to three  $\mu$ g of each plasmid is used in each 6 shot microcarrier preparation. Genes are delivered to the target tissue cells using the PDS-1000He Biolistics device. The settings on the Biolistics device are as follows: 8 mm between the rupture disc and the macrocarrier, 10 mm between the macrocarrier and the stopping screen and 7 cm between the stopping screen and the target. Each target plate is shot twice using 650 psi rupture discs. A 200 X 200 stainless steel mesh (McMaster-Carr, New Brunswick, NJ) is placed between the stopping screen and the target tissue.

Seven days after gene delivery, target tissue pieces are transferred from the high osmotic medium to selection media. For selection using the BAR gene, target tissue pieces are placed onto maintenance medium containing 100 mg/L glufosinate ammonium (Basta®) or 20 mg/L bialaphos (Herbiace®). All amino acids are removed from the selection media. After 5 to 8 weeks on these high level selection media, any growing callus is subcultured to media containing 3-20 mg/L Basta®.

For selection using the Mannose Phosphate Isomerase gene, target tissues are placed on their respective maintenance media containing no sucrose and 1% mannose. The amino acids are not removed from these media. After 5 to 8 weeks, growing callus is either subcultured to D callus maintenance medium containing no sucrose and 1.5% mannose or KM callus maintenance medium containing 1% sucrose and 0.5% mannose. Embryogenic callus growing on selection media is subcultured every 2 weeks for 4 to 8 weeks until enough callus is produced to generate 10-20 plants. Tissue surviving selection from an original target tissue piece is subcultured as a single colony and designated as an independent transformation event.

At that point, colonies selected on Basta® are transferred to a modified MS medium (Murashige and Skoog, *Physiol. Plant*, 15:473-497, 1962) containing 3% sucrose (MSS) with no selection agent and placed in the light. Either 0.25 mg/L ancymidol and 0.5 mg/L kinetin are added to this medium to induce embryo germination or 2 mg/L benzyl adenine is added. Colonies selected using mannose are transferred onto a modified MS medium containing 2% sucrose and 1% mannose (MS2S + 1M) with the ancymidol and kinetin additions described above or a modified MS medium containing 2% sucrose and 0.5% mannose (MS2S + 0.5M) with the benzyl adenine addition described above.

Regenerating colonies from Basta® selection are transferred to MS3S media without ancymidol and kinetin or benzyl adenine after 2 weeks. Regenerating colonies from mannose selection are transferred to MS2S + 1M and MS2S + 0.5M media respectively without hormones after 2 weeks. Regenerating shoots with or without roots from all colonies are transferred to Magenta boxes containing MS3S medium and small plants with roots are eventually recovered and transferred to soil in the greenhouse.

Plants are tested for expression of the PMI gene using a modified 48-well chlorophenol red assay where the media contains no sucrose and 0.5% mannose. Leaf samples (~5 mm x 5 mm) are placed on this assay media and grown in the dark for ~72 hours. If the plant is expressing the PMI gene, it can metabolize the mannose and the media will turn yellow. If not, the media will remain red.

Transformation events have also been created using Type I callus obtained from immature zygotic embryos using standard culture techniques. For gene delivery, approximately 300 mg of the Type I callus is prepared by subculturing to fresh media 1 to 2 days prior to gene delivery, selecting target tissue pieces and placing them in a ring pattern 10 mm from the center of the target plate on medium again containing 12% sucrose. After approximately 4 hours, the tissue is bombarded using the PDS-1000/He Biolistic device from DuPont. The plasmids to be transformed are precipitated onto 1 µm gold particles using the standard protocol from DuPont. Genes are delivered using two shots per target plate at 650 psi. Approximately 16 hours after gene delivery the callus is transferred to standard culture medium containing 2% sucrose with no selection agent. At 12 or 13 days after gene delivery, target tissue pieces are transferred to selection media containing 40 mg/l phosphinothricin as either Basta or bialaphos. The callus is subcultured on selection for 12 to 16 weeks, after which surviving and growing callus is transferred to standard

regeneration medium containing 3 mg/l phosphinothricin as Basta for the production of plants.

#### **Example 14: Transformation of soybean**

Protoplasts of *Glycine max* are prepared by the methods as described by Tricoli et al., 1986 (Plant Cell Rep. 5: 334-337), or Chowhury and Widholm, 1985 (Plant Cell Rep. 4: 289-292), or Klein et al., 1981 (Planta 152: 105-114). The protoplast suspension is distributed as 1 ml aliquots into plastic disposable cuvettes. For transformation, 10 µg of DNA is added in 10 µl sterile distilled water and sterilized as described by Paszkowski et al., 1984 (EMBO J. 3: 2717-2722). The solution is mixed gently and then subjected at room temperature (24 to 28°C) to a pulse of 400 Vcm<sup>-1</sup> with an exponential decay constant of 10 ms from a BTX-Transfector 300 electroporation apparatus using the 471 electrode assembly.

The above is repeated with one or more of the following modifications:

- (1) The voltage used is 200 Vcm<sup>-1</sup>, or between 100 Vcm<sup>-1</sup> and 800 Vcm<sup>-1</sup>
- (2) The exponential decay constant is 5 ms, 15 ms or 20 ms
- (3) 50 µg of sheared calf thymus DNA in 25 µl sterile water is added together with the plasmid DNA
- (4) The plasmid DNA is linearized before use by treatment with an appropriate restriction enzyme (e.g. BamHI)

The protoplasts are cultured as described in Klein et al., 1981 (Planta 152: 105-114), Chowhury and Widholm, 1985 (Plant Cell Rep. 4: 289-292), or Tricoli et al., 1986 (Plant Cell Rep. 5: 334-337), without the addition of alginate to solidify the medium.

#### **Example 15: Transformation of cotton**

*Agrobacterium* strains containing the binary vectors for transformation that are constructed by standard methods are grown 18 to 24 hours in glutamate salts media adjusted to pH 5.6 and supplemented with 0.15% mannitol, 50 µg/ml kanamycin, 50 µg/ml spectinomycin and 1 mg/ml streptomycin before they are diluted to an OD<sub>600</sub> of 0.2 in the same media without the antibiotics. The bacteria are then grown for three to five hours before dilution to an OD<sub>600</sub> of 0.2 to 0.4 and then used for inoculation of discs cut from surface sterilized cotton seeds.

The cotton seeds are soaked 20 min in 10% chlorox and rinsed with sterile water. The seeds are germinated on 0.7% water agar in the dark. The seedlings are grown for one week before inoculation of the bacteria onto the cotyledon surface.

The inoculated cotyledons are allowed to form callus before they are cut and placed on 0.7% agar containing MS salts, 3% sucrose, 100 µg/ml carbenicillin, and 100 µg/ml mefoxim. The callus is transferred to fresh media every three weeks until sufficient quantity is obtained for 4 plates. Half of the callus growing from the virulent *Agrobacterium* strains is transferred to media without hormones containing 50 µg/ml kanamycin.

#### **Example 16: *Aspergillus flavus*/Aflatoxin Disease Assay**

##### *Inoculum Production and Inoculation Protocol:*

Inoculum is an equal mixture of conidia from four highly virulent isolates of *Aspergillus flavus*. Each isolate is grown separately in petri dishes on potato-dextrose agar for 12 to 16 days at 28°C with 12h light. Cultures, including media, are blended with distilled water and filtered through double layered cheese cloth. Conidial concentrations are estimated using a hemacytometer and adjusted with distilled water. Two drops of Tween 20 per 100 ml are added as a surfactant. Conidial suspensions are prepared immediately prior to use and stored on ice while transporting from the lab to the field.

Primary ears of each plant are inoculated 20-24 days at the mid silk growth stage (50 percent of the ears with emerged silks) with a spore suspension of  $1 \times 10^6$  conidia/ml using a pin-board inoculator [*Plant Disease* (1994) 78:778-781].

##### *Ear Rot Rating and Aflatoxin Analysis:*

Forty to forty-five days after inoculation, ears are husked and a visual disease severity rating of 1 to 9 (1= no disease, 9= 90-100 percent infected) is made for each inoculated ear and averaged for each plot.

If necessary, ears are dried by forced air and shelled. Kernels for each plot are bulked and subsampled for mycotoxin analysis.

The subsamples are ground with a Romer Mill (Model 2A) and analyzed for total aflatoxin by thin layer chromatography using standard AOAC methods.

#### **Example 17: Construction of plant transformation vectors containing 5'-promoter fragments operably linked to GUS or GFP reporter genes**

To produce promoter::reporter fusions, pBSK+LOX4A (see Example 9) is used as template for the polymerase chain reaction (PCR). Gene-specific primers are used to amplify the 5' promoter region of the gene. Using combinations of the reverse primer R1 (SEQ ID NO:12) with forward primers F1 (SEQ ID NO:13) and F2 (SEQ ID NO:14) the regulatory sequences

that are ~1.2 kb and ~2 kb upstream of the initiating methionine are isolated. The nucleotide sequence of the PCR fragment amplified with forward primer F1 and reverse primer R1 is shown in SEQ ID NO:18, and the nucleotide sequence of the PCR fragment amplified with forward primer F2 and reverse primer R1 is shown in SEQ ID NO:19. For ease of cloning the primers consist of gene specific sequences and attB recombination sites for the GATEWAY™ cloning technology (Life Technologies, Invitrogen Corporation, Carlsbad, California USA). As reverse primer, primer R1 is used, which has the following sequence: 5'-CAAGAAAGCTGGGTTGACAAATTAAGTTGTCAGTGTG-3' (SEQ ID NO:12). The gene specific sequence of reverse primer R1 is underlined (corresponds to position 1356 to 1334 of SEQ ID NO:2), the attB recombination sequence is denoted in italics. Forward primers are the primers F1 and F2. Forward primer F1 has the following sequence: 5'-CAAAAAAGCAGGCTTGTAACATCCTACTCCTATTGTG-3' (SEQ ID NO:13). The gene specific sequence of forward primer F1 is underlined (corresponds to bases 159 to 181 of SEQ ID NO:2), the attB recombination sequence is denoted in italics. F1 in combination with R1 amplifies a fragment of ~1.2 kb. Forward primer F2 has the following sequence: 5'-CAAAAAAGCAGGCTCCCCGTCTTTATCTACTC-3' (SEQ ID NO:14). The gene specific sequence of forward primer F2 is underlined (corresponds to bases 31 to 48 of SEQ ID NO:1), the attB recombination sequence is denoted in italics. Primer F2 in combination with primer R1 amplifies a fragment of ~2 kb.

Using a nested PCR strategy the regulatory sequence is amplified first with primers F1+R1 or F2+R1 followed by a second PCR with primer attB1

(5'-GGGGACAAGTTTGTACAAAAAGCAGGCT-3', SEQ ID NO:15) and primer attB2 (5'-GGGGACCACTTTGTACAAGAAAGCTGGGT-3', SEQ ID NO:16). The following PCR conditions are used with gene-specific primers F1+R1 or F2+R1:

((94°C:15min):(94°C:10sec/53°C:10sec/72°C:1min)X15:(72°C:2min)). Following PCR the products are used in a second PCR reaction for amplification with the attB1+attB2 primers. In the subsequent PCR amplification, the following PCR conditions are used:

((94°C:15sec):(94°C:15sec/68°C:2min,15sec)X25:(68°C:3min). The resulting PCR product are then flanked by attB recombination sites and are used to generate Entry Clones in pDONR201 via the BP reaction according to manufacturers protocol (see: Instruction Manual of GATEWAY™ Cloning Technology, GIBCO BRL, Rockville, MD USA, <http://www.lifetech.com/>). The resulting plasmids contain ~1.2 kb and ~2 kb 5' of the *RCI-1* initiation codon and are referred to as pENTR+LOXp1.2, pENTR+LOXp2.

### 1. Produced entry vector constructs

- pENTR+LOX1.2 (pDONR201 + 1.2kb promoter fragment flanked by att recombination sequences)
- pENTR+LOX2 (pDONR201+2kb promoter fragment flanked by att recombination sequences)

These entry vectors are used to construct a binary promoter::reporter plasmid for maize or rice transformation. The regulatory/promoter sequence is fused to the GUS reporter gene (Jefferson et al, 1987, EMBO J 6: 3901-3907) or to GFP by recombination using GATEWAY™ Technology according to manufacturers protocol as described in the Instruction Manual (GATEWAY™ Cloning Technology, GIBCO BRL, Rockville, MD <http://www.lifetech.com/>). Briefly, according to this protocol the promoter fragment in the entry vector is recombined via the LR reaction with a binary *Agrobacterium* destination vector containing the GUS coding region with intron or GFP that have an attR site 5' to the GUS or GFP reporter gene (pNOV2347 or pNOV2361, respectively). The orientation of the inserted fragment is maintained by the att sequences and the final construct is verified by sequencing. The construct is then transformed into *Agrobacterium tumefaciens* strains by electroporation.

pNOV2347 and pNOV2361 are binary vectors with VS1 origin of replication, a copy of the *Agrobacterium* virG gene in the backbone and a Maize Ubiquitin promoter- PMI gene-nos terminator expression cassette between the left and right borders of T-DNA. PMI (phosphomannose isomerase) is the coding region of the *E.coli* manA gene (Joersbo and Okkels, 1996, Plant Cell Reports 16:219-221, Negrotto et al., 2000, Plant Cell Reports 19:798-803). The nos (nopaline synthase) terminator is obtained from *Agrobacterium tumefaciens* T-DNA (Depicker et al., 1982, J. Mol. Appl. Genet. 1 (6), 561-573). The maize ubiquitin promoter, the phosphomannose isomerase coding region and the nos terminator are located at nt 4114 to nt 5114, nt 6192 to nt 7295 and nt 7356 to 7604 respectively, of pNOV2347 (SEQ ID NO: 20). pNOV2361 is identical to pNOV2347, except that pNOV2361 has a GFP instead of a GUS reporter gene. The reporter-promoter cassettes are inserted closest to the right border. The selectable marker expression cassette in the binary vectors is closest to the left border. The vector contains GATEWAY™ recombination components which were introduced into the binary vector backbone by ligating a blunt-ended cassette

containing attR sites, ccdB and chloramphenicol resistance marker using the GATEWAY™ Vector Conversion System (LifeTechnologies, [www.lifetech.com](http://www.lifetech.com)). The GATEWAY™ cassette is located between nt 2351 and 4050 (complementary) of pNOV2347 and between nt 9201 and 10910 of pNOV2361. The promoter cassettes are inserted through an LR recombination reaction (LifeTechnologies, [www.lifetech.com](http://www.lifetech.com)) whereby the DNA sequence of pNOV2347 between nt 2351 and nt 4050 is removed and replaced with the LOX promoter fragment flanked by att sequences. The recombination results in the promoter sequence fused to the GFP or GUS reporter gene with intron (GIG) sequence. The GIG gene contains the ST-LS1 intron from *Solanum tuberosum* at nt 385 to nt 576 of GUS (SEQ ID NO:21) (obtained from Dr. Stanton Gelvin, Purdue University, and described in Narasimhulu, et al 1996, Plant Cell, 8: 873-886.). Shown below are the orientations of the selectable marker and promoter-reporter cassettes in the binary vector constructs.

## 2. Produced constructs for stable transformation

- pNOV6800 (RB nos + GIG gene + LOX1.2 promoter fragment – ZmUbi + PMI gene + nos LB)
- pNOV6801 (RB LOX1.2 promoter fragment + GFP gene + nos-ZmUbi + PMI gene + nos LB)

The nucleotide sequence of pNOV6800 is depicted in SEQ ID NO:22. pNOV6800 and pNOV6801 differ only in the expression cassette located between the right and left borders of the binary vector.

## 3. Constructs used for comparison

- pNOV2110 (RB ZmUbi Promoter + GFP gene + nos-ZmUbi + PMI gene + nos LB)
  - pNOV 3640 (RB nos-GIG-ZmUbi promoter nos-AtPPOdm-ZmUbi promoter LB)
- GUS – Intron - GUS, GFP and polyA fragments are identical to those used for the LOX promoter constructs above. The ZmUbi promoter corresponds to the fragment from base 12 to base 2009 in pNOV2110 and contains promoter, Exon1 and Intron1 of the Maize Ubiquitin gene. The AtPPOdm sequence encodes a mutated form of the protophorinogen oxidase protein which confers resistance to herbicides (PPO inhibitors) that normally inactivate the enzyme (US patent no. 5,939,602).



**Example 18: *Agrobacterium*-mediated transformation of Maiz**

Transformation of immature maize embryos is performed essentially as described in Negrotto et al., (2000) Plant Cell Reports 19: 798-803. For this example, all media constituents are as described in Negrotto et al., *supra*. However, various media constituents described in the literature may be substituted.

**1. Transformation plasmids and selectable marker**

The genes used for transformation are cloned into a vector suitable for maize transformation as described in Example 17. Vectors used contain the phosphomannose isomerase (PMI) gene (Negrotto et al. (2000) Plant Cell Reports 19: 798-803) as a selectable marker.

**2. Preparation of *Agrobacterium tumefaciens***

*Agrobacterium* strain LBA4404 (pSB1) containing the plant transformation plasmid is grown on YEP (yeast extract (5 g/L), peptone (10g/L), NaCl (5g/L), 15g/l agar, pH 6.8) solid medium for 2 to 4 days at 28°C. Approximately  $0.8 \times 10^9$  *Agrobacteria* are suspended in LS-inf media supplemented with 100 µM acetosyringone (As) (LSAs medium) (Negrotto et al., (2000) Plant Cell Rep 19: 798-803). Bacteria are pre-induced in this medium for 30-60 minutes.

**3. Inoculation**

Immature embryos from A188 or other suitable maize genotypes are excised from 8 – 12 day old ears into liquid LS-inf + 100 µM As (LSAs). Embryos are rinsed once with fresh infection medium. *Agrobacterium* solution is then added and embryos are vortexed for 30 seconds and allowed to settle with the bacteria for 5 minutes. The embryos are then transferred scutellum side up to LSAs medium and cultured in the dark for two to three days. Subsequently, between 20 and 25 embryos per petri plate are transferred to LSDc medium supplemented with cefotaxime (250 mg/l) and silver nitrate (1.6 mg/l) (Negrotto et al. 2000) and cultured in the dark for 28°C for 10 days.

**4. Selection of transformed cells and regeneration of transformed plants**

Immature embryos producing embryogenic callus are transferred to LSD1M0.5S medium (LSDc with 0.5 mg/l 2,4-D instead of Dicamba, 10g/l mannose, 5 g/l sucrose and no silver

nitrate). The cultures are selected on this medium for 6 weeks with a subculture step at 3 weeks. Surviving calli are transferred either to LSD1M0.5S medium to be bulked-up or to Reg1 medium (as described in Negrotto et al., 2000). Following culturing in the light (16 hour light/ 8 hour dark regiment), green tissues are then transferred to Reg2 medium without growth regulators (as described in Negrotto et al. 2000) and incubated for 1-2 weeks. Plantlets are transferred to Magenta GA-7 boxes (Magenta Corp, Chicago Ill.) containing Reg3 medium (as described in Negrotto et al. 2000) and grown in the light. Plants that are PCR positive for the promoter-reporter cassette are transferred to soil and grown in the greenhouse.

#### **EXAMPLE 19: GUS Reporter gene assays**

Promoter activity is evaluated qualitatively and quantitatively using histochemical and florescence assays for expression of the  $\beta$ -glucuronidase (GUS) enzyme.

##### 1. Histochemical $\beta$ -glucuronidase (GUS) assay

For qualitative evaluation of promoter activity, various tissues and organs are used in GUS histochemical assays. Either whole organs or pieces of tissue are dipped into GUS staining solution. GUS staining solution contains 1 mM 5-bromo-4-chloro-3-indolyl glucuronide (X-Gluc, Duchefa, 20 mM stock in DMSO), 100 mM Na-phosphate buffer pH 7.0, 10 mM EDTA pH 8.0, and 0.1% Triton X100. Tissue samples are incubated at 37°C for 1-16 hours. If necessary samples can be cleared with several washes of 70% EtOH to remove chlorophyll. Following staining tissues are viewed under a light microscope to evaluate the blue staining showing the GUS expression pattern.

##### 2. $\beta$ -glucuronidase (GUS) fluorescence assay

For quantitative analysis of promoter activity in various tissues and organs, GUS expression is measured fluorometrically. Tissue samples are harvested and ground in ice cold GUS extraction buffer (50mM Na<sub>2</sub>HPO<sub>4</sub> pH 7.0, 5mM DTT, 1mM Na<sub>2</sub>EDTA, 0.1% TritonX100, 0.1% sarcosyl). Ground samples are spun in a microfuge at 10,000 rpm for 15 minutes at 4 °C. Following centrifugation the supernatant is removed for GUS assay and for protein concentration determination.

To measure GUS activity the plant extract is assayed in GUS assay buffer (50mM Na<sub>2</sub>HPO<sub>4</sub> pH 7.0, 5mM DTT, 1mM Na<sub>2</sub>EDTA, 0.1% TritonX100, 0.1% sarcosyl, 1mM 4-

Methylumbelliferyl-beta-D-glucuronic acid dihydrate (MUG)), prewarmed to 37 °C. Reactions are incubated and 100 µL aliquots are removed at 10 minute intervals for 30 minutes to stop the reaction by adding to tubes containing 900 µL of 2% Na<sub>2</sub>CO<sub>3</sub>. The stopped reactions are then read on a Tecan Spectrofluorometer at 365 nm excitation and 455 nm emission wavelengths. Protein concentrations are determined using the BCA assay following manufacturers protocol. GUS activity is expressed as relative fluorometric units (RFU)/mg protein.

#### **EXAMPLE 20: GFP Reporter gene assays**

Promoter activity is evaluated qualitatively using microscopic imaging fluorescence and quantitatively using fluorescence assays for expression of the green fluorescent protein.

##### 1. Microscopic evaluation of GFP expression

Expression of the promoter::GFP fusion is monitored in transformants by microscopic imaging using a Leica MzFLIII fluorescence microscope (Leica Microsystems, Heidelberg, Germany) with GFP2 and GFP3 filter settings.

##### 2. Quantitative GFP Fluorometric Assay

To assay expression of GFP in tissues of transgenic plants, harvested tissue is frozen and frozen tissue is ground thoroughly. Following grinding add 300 µL of extraction buffer (EB; 10mM Tris-HCL (pH7.5), 100 mM NaCl, 1mM MgCl<sub>2</sub>, 10mM DTT and 0.1% Sarcosyl). Vortex well to mix sample, centrifuge for 10 minutes, then transfer the supernatant to a new tube or microtitre plate well for reading. The sample tube or plate is then inserted into the Tecan Spectrofluorometer plate reader set to 465 nm excitation and 512 nm emissions wavelengths for 10 flashes and a gain of 100. If the expression levels are too high resulting in some of the samples being over the limit (says VALUE in the cell) then parameters are changed to 8 flashes and a gain of 80.

Protein concentrations are determined using the BCA assay following manufacturers protocol. GFP activity is expressed as the relative fluorescence units (RFU) per milligram of protein.

**Example 21: Evaluation of promoter activity**

To evaluate activity of the promoter::reporter fusion, tissue samples are harvested from untreated control plants as well as from plants treated with the chemical activator, BTH or jasmonic acid. In transgenic rice treatment with the chemical inducers is done 10 days after sowing T1 seed at the emergence of leaf 3. In transgenic maize treatment with the chemical inducers is done three weeks after sowing T1 seed. All chemical concentrations are given as ppm (mg active ingredient l<sup>-1</sup> of applied solution). Probenazole is applied as a 250 ppm solution of the pure substance by soil drench as described (Thieron et al. (1995) Systemic acquired resistance in rice: Studies on the mode of action of diverse substances inducing resistance in rice to *Pyricularia oryzae*. *Mededelingen Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen Universiteit Gent*. 60, 421-430). Formulations of BTH (1:1 (w/w) mixture of active ingredient and wettable powder) is applied onto leaves by spraying. All controls are done by application of spray-solutions without active substance. Jasmonic acid is applied as a 1 mM solution in ethanol as described (Schweizer et al. (1997) *Plant Physiol.* 114, 79-88). Wounding and measurement of gene expression in systemic tissue is done according to (Schweizer et al. (1998) *Plant J.* 14, 475-481).

Various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and examples. Such modifications are intended to fall within the scope of the appended claims.

Various references and patents have been cited herein, and are all incorporated by reference in their entireties.

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>28</u> , line <u>10-19</u>	
<b>B. IDENTIFICATION OF DEPOSIT</b> <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution <div style="text-align: center;"><b>DEUTSCHE SAMMLUNG VON MIKROORGANISMEN UND ZELLKULTUREN GmbH (DSMZ)</b></div>	
Address of depositary institution (including postal code and country)  <div style="text-align: center;">Mascheroder Weg 1b D-38124 Braunschweig Germany</div>	
Date of deposit <div style="text-align: center;">06 June 2000 (06.06.00)</div>	Accession Number <div style="text-align: center;">DSM 13524</div>
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
We request the Expert Solution where available.	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indications are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<div style="text-align: center; border-bottom: 1px solid black; margin-bottom: 5px;">For receiving Office use only</div> <div style="border: 1px solid black; padding: 5px;"> <input type="checkbox"/> This sheet was received with the international application       </div> <div style="border: 1px solid black; padding: 5px; margin-top: 5px;">         Authorized officer       </div>	<div style="text-align: center; border-bottom: 1px solid black; margin-bottom: 5px;">For International Bureau use only</div> <div style="border: 1px solid black; padding: 5px;"> <input type="checkbox"/> This sheet was received by the International Bureau on:       </div> <div style="border: 1px solid black; padding: 5px; margin-top: 5px;">         Authorized officer       </div>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>28</u> , line <u>10-19</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution <b>Agricultural Research Service Culture Collection (NRRL) National Center for Agricultural Utilization Research Agricultural Research Service, United States Department of Agriculture</b>	
Address of depositary institution (including postal code and country)  <b>1815 North University Street Peoria, Illinois 61604 United States of America (USA)</b>	
Date of deposit <b>25 May 2001 (25.05.01)</b>	Accession Number <b>NRRL B-30480</b>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
We request the Expert Solution where available.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	
<input type="checkbox"/> This sheet was received with the international application	
Authorized officer	

For International Bureau use only	
<input type="checkbox"/> This sheet was received by the International Bureau on:	
Authorized officer	

***What is claimed is:***

1. An isolated nucleic acid molecule capable of driving chemically-inducible but not wound- or pathogen-inducible expression of an associated nucleotide sequence.
2. The isolated nucleic acid molecule according to claim 1, wherein said nucleic acid molecule is a component of the *Pst*I/*Pst*I fragment of about 4.5 kb in length from plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524.
3. The isolated nucleic acid molecule according to claim 1, wherein said nucleic acid molecule is a component of the nucleotide sequence depicted in SEQ ID NO:17.
4. The isolated nucleic acid molecule according to claim 3, wherein said nucleic acid molecule is depicted in SEQ ID NO:18.
5. The isolated nucleic acid molecule according to claim 3, wherein said nucleic acid molecule is depicted in SEQ ID NO:19.
6. The isolated nucleic acid molecule according to claim 1, wherein said nucleic acid molecule comprises the nucleotide sequence depicted in SEQ ID NO:1.
7. The isolated nucleic acid molecule according to claim 1, wherein said nucleic acid molecule comprises nt 1 to nt 1358 of the nucleotide sequence depicted in SEQ ID NO:2.
8. The isolated nucleic acid molecule according to claim 1, wherein said nucleic acid molecule comprises nt 1702 to nt 2104 of SEQ ID NO:2 and/or nt 1 to nt 97 of SEQ ID NO:3 and/or nt 367 to nt 1283 of SEQ ID NO:3 of SEQ ID NO:3.
9. The isolated nucleic acid molecule according to claim 1 or 2, wherein said nucleic acid molecule comprises a combination of any one of the nucleotide sequences or portions thereof depicted in SEQ ID NO:1, SEQ ID NO:2 and SEQ ID NO:3.
10. An isolated nucleic acid molecule which hybridizes under stringent conditions to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or

to the 4.5 kb *Pst*I fragment of plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524, wherein said isolated nucleic acid molecule is capable of driving chemically-inducible but not wound- or pathogen-inducible expression of an operably-linked nucleotide sequence.

11. An isolated nucleic acid molecule comprising a consecutive stretch of at least 50 nt of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or of the 4.5 kb *Pst*I fragment of plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524, wherein said isolated nucleic acid molecule is capable of driving chemically inducible but not wound- or pathogen-inducible expression of an operably-linked nucleotide sequence.

12. An isolated nucleic acid molecule according to claim 11 wherein said consecutive stretch of at least 50 nt has at least 70% sequence identity with a consecutive stretch of at least 50 nt of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19, or of the 4.5 kb *Pst*I fragment of plasmid pBSK+LOX4A which has been deposited under accession no DSM 13524.

13. The isolated nucleic acid molecule according to any one of claims 1 to 12, wherein the chemical inducer is selected from the group consisting of BTH (benzo(1,2,3)thiadiazole-7-carbothioic acid S-methyl ester), INA (2,6-dichloroisonicotinic acid) and probenazole.

14. The isolated nucleic acid molecule according to any one of claims 1 to 12, wherein the chemical inducer is jasmonic acid.

15. A recombinant nucleic acid molecule comprising the isolated nucleic acid molecule according to any one of claims 1 to 14 operably linked to a nucleotide sequence of interest.

16. The recombinant nucleic acid molecule of claim 15, wherein the nucleotide sequence of interest comprises a polypeptide coding sequence.

17. The recombinant nucleic acid molecule of claim 16, wherein the coding sequence comprises at its 5'-end a nucleotide sequence encoding the amino acid sequence depicted in SEQ ID NO:6.



18. The recombinant nucleic acid molecule of claim 16 or 17, wherein the coding sequence encodes a desirable phenotypic trait.
19. The recombinant nucleic acid molecule of claim 16, wherein the coding sequence is in antisense orientation.
20. A nucleic acid expression vector comprising the isolated nucleic acid molecule according to any one of claims 1 to 14 or the recombinant nucleic acid molecule of any one of claims 15 to 19.
21. A host cell stably transformed with an isolated nucleic acid molecule according to any one of claims 1 to 14 or a recombinant nucleic acid molecule of any one of claims 15 to 19.
22. The host cell of claim 21, wherein said host cell is a plant cell.
23. A plant and the progeny thereof stably transformed with an isolated nucleic acid molecule according to any one of claims 1 to 14 or a recombinant nucleic acid molecule of claims 15 to 19.
24. The plant of claim 23, wherein said plant is selected from the group consisting of maize, wheat, sorghum, rye, oats, turf grass, rice, barley, soybean, cotton, tobacco, sugar beet and oilseed rape.
25. Use of the nucleic acid molecule of any one of claims 1 to 14 to express a nucleotide sequence of interest.
26. A method of producing the isolated nucleic acid molecule according to claim 1, wherein the isolated nucleic acid molecule is produced by a polymerase chain reaction wherein at least one oligonucleotide used comprises a sequence of nucleotides which represents a consecutive stretch of 15 or more nucleotides of SEQ ID NO:1, SEQ ID NO:2 SEQ ID NO:3, SEQ ID NO:17, SEQ ID NO:18 or SEQ ID NO:19.

27. An isolated nucleic acid molecule encoding the amino acid sequence depicted in SEQ ID NO:6, wherein said amino acid sequence is capable of targeting an associated protein to plastids.
28. The isolated nucleic acid molecule according to claim 27, wherein said isolated nucleic acid molecule is depicted in SEQ ID NO:4.
29. An isolated nucleic acid molecule which hybridizes under stringent conditions to the nucleotide sequence depicted in SEQ ID NO:4.
30. A peptide encoded by the nucleotide sequence of any one of claims 27 to 29.
31. A peptide encoded by the nucleotide sequence of claim 28.
32. Use of the peptide of claim 30 or 31 to target an associated protein of interest to plastids.
33. An isolated nucleic acid molecule which hybridizes under stringent conditions to SEQ ID NO:5, wherein the protein encoded by said DNA has at least 65% amino acid sequence identity with the amino acid sequence depicted in SEQ ID NO:7 and encodes a protein with lipoxygenase activity.
34. The isolated nucleic acid molecule according to claim 33, wherein said nucleotide sequence encodes the protein depicted in SEQ ID NO:7.
35. A protein encoded by the nucleotide sequence of the isolated nucleic acid molecule of claim 33 or 34.
36. A protein encoded by the nucleotide sequence of the isolated nucleic acid molecule of claim 34.
37. Use of the protein according to claim 35 or 36 to inhibit fungal mycotoxins.

- 58 -

38. A method of inhibiting fungal mycotoxins in a plant by transforming the plant with nucleic acid molecule of claim 33 or 34 and expressing a polypeptide having lipoxygenase activity.

39. A recombinant nucleic acid molecule comprising the nucleic acid molecule according to any one of claims 27 to 29 or claims 33 to 34.

40. A host cell stably transformed with the recombinant nucleic acid molecule of claim 39.

41. The host cell of claim 40, wherein said host cell is a plant cell.

42. A plant and the progeny thereof stably transformed with the recombinant nucleic acid molecule of claim 39.

43. The plant of claim 42, wherein said plant is selected from the group consisting of maize, wheat, sorghum, rye, oats, turf grass, rice, barley, soybean, cotton, tobacco, sugar beet and oilseed rape.

44. A seed of the plant and progeny thereof of claim 43.

- 1 -

## SEQUENCE LISTING

<110> Syngenta Participations AG  
University of Zurich

<120> Lipoyxygenase Genes, Promoters, Transit Peptides and Proteins Thereof

<130> A-31484A

<140>

<141>

<150> GB 0017275.9

<151> 2000-07-13

<150> GB 0022739.7

<151> 2000-09-15

<160> 22

<170> PatentIn Ver. 2.1

<210> 1

<211> 358

<212> DNA

<213> Oryza sativa

<400> 1

```
ctgcaggccc agcaaacacg cgtcacacgt ccccgctctt atctactcgc gcccatcctc 60
tacttatcta ctggcctct cctcctccac tactcctcgc ctggtcctcc cctctcctcg 120
atccctctt ccccttcccg ggctctccg tgggtggttg cgcgcggcgg caggcggagg 180
cgggcgggtg tgcgcggcgg cgtgacataa gcggcgggct gatgggaggg cgggcggcgg 240
gtggagggga ggatggtggc gggcggatgg gtgggcgaca acgggtgggc ggcagcggga 300
ggcggtgccc tcccttctgc cggaggggag gcgggaggcg gcagcggcag cggcggat 358
```

<210> 2

<211> 2104

<212> DNA

<213> Oryza sativa

<400> 2

```
actatgtttg tcggagtgtg tacatagata gggggccatt cgatgggacc gctttctatg 60
agttcggggg tgtcgcacgt gagggagcgc tcgggttccc attgaggacg gaaacaaatg 120
aacaacaat aattatatta gtttgagaca tccggatgtg taacatccta ctctattgt 180
gtcatgtgtt ttctctcttg ctagtgaatg atcttatatg tcagacttag agtcgtagtt 240
catctctttt ggatttccga agaacttcac tgtgtatggc tatatgttct aataactctg 300
tatgttcgaa cctcctttct cttggtcatg gctctcccat tatactatc tagggacacc 360
acatgatcag aatattaggg agatgctaata taagtgcata gtgccggatt atgtaaggaa 420
aatcttgtca gcactgctcc tctctccctt tttgcgaggc ttagcgtaga accacgaaaa 480
aaaaacttgt attaaagaca ccaagataac tagctgcaaa catcccagac aagccgatca 540
tactcgaaat caaccacctt ctctatcaca tcatattgaa tttaaggttt atccaatcac 600
attcaaagt actaactaca gcaaccagga aaatcctcga aagaattaga agcatcaatg 660
aagaagcatt gaacatcact tctataatcc ctccaggaaa ccatcagaag taaaacattc 720
tctggccccc attggttgaa tttttttcta ataagccaaa acggcttatt agagaataaa 780
```

- 2 -

```

aataaatgtg taggtaaaac ttttatatat gtgttttttt taacttaaaa gccaatactg 840
aaaaaaacta cgttgaaaa atctcagaat caatctcaaa attaagtttg aaaattcaaa 900
at ttgtggctta ttctttggct tattgggcca tctgatggga gcctctatat ttggtagcac 960
ataactaaat aaataaaaagg gttcatattc ctaattgtcg aaagtctctc aaagtgcacat 1020
aacacactag tatttacact acggacaacc ctccctagttt agacttttaga gttacatgtt 1080
gccagtttc gaggtaaagt agactagctc aaatgtgtct caaacagtcg ttcccggaa 1140
ataagtcaca agtggtccca cgtgtaggca tgtttcacgc ttagatcgat cgagtttcgt 1200
ttocatctgt acgtacgttg taccaccca acccgatgat cgatatgatc gatcgcccc 1260
tataaattgg acatcctgga catgctcttt cttcagacat ctccaccctt cctatcttga 1320
tatataactt cgacacactg acaacttaat ttgtcagaat gctcacggcc acgcagactc 1380
tggtcgccggc agtgctctcc cggagccatg gcgccccttc ttcttctcc agccagccgc 1440
gcgcacccgc cgccgcggcc tcgagagtaa gctgcaccgc cgtcggcgcc ttgtcggagg 1500
tcgtcaatgg cgaactcgtc gtcggcgacc aagaacagac gaccgacgac ctccctacgc 1560
ggcacaagaa tgcgtcgcc gactacacgc tgagcgccac ggtgacggtg agcttgaagc 1620
aggacgattc cactccccag aaggtggcgg acatggttaa tcgagactgg cttttccttg 1680
at ttcttcag ctgcataata ggtgatgaca agctccatag ctactctatc gatcgtagcg 1740
gcgcctctt tcgcgccata tataatcttg aacatgggca ctgattccca ttcgaaaaaa 1800
tatcttaatt tgagaggatg ctttctactg gtagctagca ttctactact agaattatcg 1860
tattaaaagt gatcgagcc gttcgattta ctagctagct cgaatgatag catatattgt 1920
aagtgtaaat tttgatacat tttgatatat aacttcgaca ttaggaaaaa aactacatta 1980
aaaatttgg tacctaaaaa actataaaat ttctactatt atcagtaaaa atattttttg 2040
cttaccatta gtagattaga ttgtatataa gatatactac ctctgttttt taatagatga 2100
cgcc 2104

```

&lt;210&gt; 3

&lt;211&gt; 1516

&lt;212&gt; DNA

<213> *Oryza sativa*

&lt;400&gt; 3

```

gtcatctatt aaaaaacgga gagagtatat gattcatcgg aattaaaaaa tagacgggtat 60
aacacattgt aaaacaccta ttttcgtcac tctacagagg ggatgcacac ggagcctcag 120
ctcgccaggt actgcacat ggatggcaaa ggctccttca tatacagggc cagcttcagc 180
atcccgctct cgttggacgc cgtcggcgcc gtgcaggctg tgaaccgcta cagcagcgag 240
gtgtacatct cggacatcga cgtccacctc tgcggcggcc gccatcagtg gaccgacatc 300
actttccact gcaactcttg gatcgactac aaccccaacg accagcgctt cttcttccct 360
ctcaaggcca ggggttcacg ttattgcca taccgatatt gttttattcg tgaggatgggt 420
aaaattagct atcgccaaa aattcctttt ttttttgaat tagctcatat gtaaaataaa 480
attaacacct gcaaaccatc gaaaagggtg gaggattata tttgcttaat tagggcagtc 540
ccaaccata acactagaca tggtttatat aacactagac atcatcaaga aactagtact 600
acacactact cttccaatgc aaataccact attccatact tcagttaa at gctacttacc 660
tcacatgatg tcttagatgt tgtgtagaaa ccatgtctca tgcaagacat ggtttccttc 720
tctttcttta tttattcact tgtcacatta tattttgtc ctaggtggca acttattta 780
tgctatggac accattctag ttattgggtt gggaatgcc ttaagggtgt gtttgattgg 840
gtggctgaag ctgcatggga gttggggata aggcagccat ctctcactag attggatgag 900
tggtatgagg gaacacatg tgggtgatgt tgctgattgg ctgtgatagg tttaacttga 960
tggttaacaat ctcaaaactga tttttttt taaactgatt atcctattta taatccgatt 1020
atatgattat atttatatta tagttaatct tcaaaacaaa atttcacatg gttatacgaa 1080
tggtgcccac tgacttattg cgtccatgct atatcctcca gcctatggat atatcaaaca 1140
aaacatagat ggggttatccc atccccatcc acttatgacc gtgaactaaa cgcgccctaa 1200
tctccatttc cgctcgatcg atcgttgttc agctgcaaat cattcacact gcttgattat 1260
tgcttgcatg caogatttg cagtcgtacc tccgctctca gacgcccagg ggctgaaga 1320
atctgcgcaa ggaagagctc agggccatcc gcggcgatgg ccgcgcgag cgcaaggagt 1380
gggagcgcat ctacgactac gacgtctaca acgacctcgg cgaccccgac aatgaccggg 1440

```

- 3 -

ccactcgctcg gccggtgctc ggcggccgcg ggcgcccta cccgcgccgc tgccgcacgg 1500  
 gccgcgccgc ctgcag 1516

&lt;210&gt; 4

&lt;211&gt; 158

&lt;212&gt; DNA

<213> *Oryza sativa*

&lt;400&gt; 4

ctatcttgat atataacttc gacacactga caacttaatt tgtcagaatg ctcacggcca 60  
 cgcagactct ggcgcggca gtgctctccc ggagccatgg cgcctctct tcttctcca 120  
 gccagccgcg ccgcaccgcc gccgcgcct cgagagta 158

&lt;210&gt; 5

&lt;211&gt; 3018

&lt;212&gt; DNA

<213> *Oryza sativa*

&lt;400&gt; 5

ctatcttgat atataacttc gacacactga caacttaatt tgtcagaatg ctcacggcca 60  
 cgcagactct ggcgcggca gtgctctccc ggagccatgg cgcctctct tcttctcca 120  
 gccagccgcg ccgcaccgcc gccgcgcct cgagagtaag ctgcaccgcg gtcggcgct 180  
 tgtcggaggt cgtcaatggc gaactcgctc tcggcgacca agaacagacg accgacgacc 240  
 tctttacgcg gcacaagaat gtcgtcgccg actacacgct gagcgccacg gtgacgggtga 300  
 gcttgaagca ggacgattcc actccccaga aggtggcgga catggttaat cgagactggc 360  
 ttttcttga tttcttcagc tcgcatatag aggggatgca cacggagcct cagctcgcca 420  
 ggtactcgca catggatggc aaaggtcct tcatatacga ggccagcttc agcatcccgt 480  
 cctcgttgga cgcgctcgcc gccgtgcagg tcgtgaaccg ctacagcagc gaggtgtaca 540  
 tctcggacat cgacgtccac ctctgcccgc gccgccatca gtggaccgac atcactttcc 600  
 actgcaactc ttggatcgac tacaaccca acgaccagcg cttcttcttc cctctcaagt 660  
 cgtacctccc gtctcagacg cccaggggcg tgaagaatct gcgcaaggaa gagctcaggg 720  
 ccatccgcg cgatggccgc ggcgagcgca aggagtggga gcgcatctac gactacgacg 780  
 tctacaacga cctcggcgac cccgacaatg acccgccac tcgtcggccg gtgctcggcg 840  
 gccgcgggcg cccctaccgc cgcgcgtgcc gcacggggcg cgcgcgtgc aggacagacc 900  
 cgtcgtcgga gtgcgcgcg gccaaaggac gcgcgggat ctacgtgcca cgggacgagg 960  
 cgttcacgga gcggaaggcc ggcgcgttcg ccaccaagaa ggcgctgtcg gcgctgtcg 1020  
 cgttcaccac ggcgcagag gtgtcggcg accggcgcg gggcttcccg tcgctggcg 1080  
 ccatcgacgc gctgtacgag gacgggtaca agaaccggcc gtcgtcgtcg cagcaggagg 1140  
 cggacaacct cgaaggctac ttcaggaggg tgcctcagaa gcagggtgaag ctgctgctca 1200  
 agggcgagaa ggaggagttc aaggaggagc tacgcaaagt gttcaaattc caaacgccg 1260  
 agattcacga caaggacaag cttgcatggc tcagagacga ggagttcgcg cggcaaacgc 1320  
 tggcagggat gaacctctc agcatccaac ttgtcaggga cacggacttc cctatatcca 1380  
 gcaagctgga cgaggaaacc tacggcccag gggactccct catcacaaa gagctgattg 1440  
 aagagcagat taatggggtc atgacagcag aggaggccgt ggagaagaag aagctgttca 1500  
 tgctggacta ccacgacgtg ctctgcccgt tcgtgcacgc ggtgcgcgag ctggacgaca 1560  
 ccacgctgta cgcctcgcg acgctcttct tctgacgga ggacggcacg ctgaggccga 1620  
 tcgccatcga gctgacgagg cccaagtccc ccaacacgcc gcagtggcg caggcttcca 1680  
 cgcggggtc cagcgtcgcg gcgtcctggc tgtggcagct cgcctaacg cagctcctcg 1740  
 cccacgacac cggctaccac cagctcgtca gccactggct gaggacgcac tgctgcgtgg 1800  
 agcgtacgt gatcgcgcg aaccggcggc tgagccagat gcacccatc taccgactgc 1860  
 tgcaccgca cttcgcctc accatggaga tcaacgccc agcgcgggg atgctcatca 1920  
 acgccaatgg aatcctcgag agcgccttcg cgcggggaa gctctgcatg gagtcagct 1980  
 cggcggttta cgacaagttt tggaggttcg acatggaggc tctgcccgc gatctcatcc 2040

- 4 -

```

ggaggggcat ggcgatcgaa tgcgaggatg gcaagctgga gctgacgata gaggactacc 2100
cgtacgccaa cgacggcctg ctcatctggg actccatcaa ggagtgggtg tcggattatg 2160
tgaaccatta ctaccagttg gcttcagaca tccacatgga caaggagctc cagggttggt 2220
ggaacgaggt gcgaaccaag ggccacccgg acaaggagga agggaggcca gagctgaact 2280
gccacgggag cctcgtcgag gttctgacca ccatcatctg ggtcgcgtcg gggcaccatg 2340
cggcggtgaa ctttggccag taccctacg ccggctactt cccaatcgc cccaccatcg 2400
cccggcgga catgccgacg gaggggcagg cgtgcagtca cgacggcatg cagccaacgt 2460
tcgttgagga tcccgtcagg gtgctactag acacgttccc atcgcagtac cagaccaccc 2520
tcgtcctgcc ggtgctcaac ctgctatcgt cacactcgcc cggcgaggag tacatgggca 2580
cgcgtgcgga gtcagcgtgg atggcgga gggaggtcag ggcggcggtc gggaggttca 2640
acgagaggat gatgagcatc ggggagatga tcgactgccg gaacaaggat ccggagcgaa 2700
agaaccggca gggccccggc gtggtgccgt acgtgctgct caagccgtcc tacggtgacc 2760
ctaaggacat gacgtccgtg atggagatgg gtatcccaaa cagcatctca atttgagttg 2820
tgccaatgag cttgcatctg tttggcgtgc tcacgtgac attatgtatg aaataaaatg 2880
gattaaaaat ccggcctcgt caaggaatgg ctaacacagc gagcctgcat ctgtttggag 2940
tgctcatcgt gcgattatga aataaaatga cctggcatct gtttgccact gttttcttgt 3000
aaaaaaaaa aaaaaaaaaa                                     3018

```

&lt;210&gt; 6

&lt;211&gt; 37

&lt;212&gt; PRT

<213> *Oryza sativa*

&lt;400&gt; 6

```

Met Leu Thr Ala Thr Gln Thr Leu Ala Pro Ala Val Leu Ser Arg Ser
 1             5             10             15

```

```

His Gly Ala Pro Ser Ser Phe Ser Ser Gln Pro Arg Arg Thr Ala Ala
      20             25             30

```

```

Ala Ala Ser Arg Val
      35

```

&lt;210&gt; 7

&lt;211&gt; 922

&lt;212&gt; PRT

<213> *Oryza sativa*

&lt;400&gt; 7

```

Met Leu Thr Ala Thr Gln Thr Leu Ala Pro Ala Val Leu Ser Arg Ser
 1             5             10             15

```

```

His Gly Ala Pro Ser Ser Phe Ser Ser Gln Pro Arg Arg Thr Ala Ala
      20             25             30

```

```

Ala Ala Ser Arg Val Ser Cys Thr Arg Val Gly Ala Leu Ser Glu Val
      35             40             45

```

```

Val Asn Gly Glu Leu Val Val Gly Asp Gln Glu Gln Thr Thr Asp Asp
      50             55             60

```

```

Leu Leu Thr Arg His Lys Asn Val Val Ala Asp Tyr Thr Leu Ser Ala

```

- 5 -

65		70		75		80
Thr Val Thr Val Ser Leu Lys Gln Asp Asp Ser Thr Pro Gln Lys Val						
	85		90			95
Ala Asp Met Val Asn Arg Asp Trp Leu Phe Leu Asp Phe Phe Ser Ser						
	100		105			110
His Ile Glu Gly Met His Thr Glu Pro Gln Leu Ala Arg Tyr Ser His						
	115		120			125
Met Asp Gly Lys Gly Ser Phe Ile Tyr Glu Ala Ser Phe Ser Ile Pro						
	130		135			140
Ser Ser Leu Asp Ala Val Gly Ala Val Gln Val Val Asn Arg Tyr Ser						
	145		150			155
Ser Glu Val Tyr Ile Ser Asp Ile Asp Val His Leu Cys Gly Gly Arg						
		165		170		175
His Gln Trp Thr Asp Ile Thr Phe His Cys Asn Ser Trp Ile Asp Tyr						
		180		185		190
Asn Pro Asn Asp Gln Arg Phe Phe Phe Pro Leu Lys Ser Tyr Leu Pro						
		195		200		205
Ser Gln Thr Pro Arg Gly Val Lys Asn Leu Arg Lys Glu Glu Leu Arg						
		210		215		220
Ala Ile Arg Gly Asp Gly Arg Gly Glu Arg Lys Glu Trp Glu Arg Ile						
		225		230		235
Tyr Asp Tyr Asp Val Tyr Asn Asp Leu Gly Asp Pro Asp Asn Asp Pro						
		245		250		255
Ala Thr Arg Arg Pro Val Leu Gly Gly Arg Gly Arg Pro Tyr Pro Arg						
		260		265		270
Arg Cys Arg Thr Gly Arg Arg Arg Cys Arg Thr Asp Pro Ser Ser Glu						
		275		280		285
Ser Pro Pro Ala Lys Asp Gly Ala Gly Ile Tyr Val Pro Arg Asp Glu						
		290		295		300
Ala Phe Thr Glu Arg Lys Ala Gly Ala Phe Ala Thr Lys Lys Ala Leu						
		305		310		315
Ser Ala Leu Ser Ala Phe Thr Thr Ala Gln Arg Val Ser Gly Asp Arg						
		325		330		335
Arg Arg Gly Phe Pro Ser Leu Ala Ala Ile Asp Ala Leu Tyr Glu Asp						
		340		345		350
Gly Tyr Lys Asn Arg Pro Ser Ser Ser Gln Gln Glu Ala Asp Asn Leu						
		355		360		365



- 6 -

Glu Gly Tyr Phe Arg Glu Val Leu Gln Lys Gln Val Lys Leu Leu Leu  
 370 375 380

Lys Gly Glu Lys Glu Glu Phe Lys Glu Glu Leu Arg Lys Val Phe Lys  
 385 390 395 400

Phe Gln Thr Pro Glu Ile His Asp Lys Asp Lys Leu Ala Trp Phe Arg  
 405 410 415

Asp Glu Glu Phe Ala Arg Gln Thr Leu Ala Gly Met Asn Pro Leu Ser  
 420 425 430

Ile Gln Leu Val Arg Asp Thr Asp Phe Pro Ile Phe Ser Lys Leu Asp  
 435 440 445

Glu Glu Thr Tyr Gly Pro Gly Asp Ser Leu Ile Thr Lys Glu Leu Ile  
 450 455 460

Glu Glu Gln Ile Asn Gly Val Met Thr Ala Glu Glu Ala Val Glu Lys  
 465 470 475 480

Lys Lys Leu Phe Met Leu Asp Tyr His Asp Val Leu Leu Pro Phe Val  
 485 490 495

His Ala Val Arg Glu Leu Asp Asp Thr Thr Leu Tyr Ala Ser Arg Thr  
 500 505 510

Leu Phe Phe Leu Thr Glu Asp Gly Thr Leu Arg Pro Ile Ala Ile Glu  
 515 520 525

Leu Thr Arg Pro Lys Ser Pro Asn Thr Pro Gln Trp Arg Gln Val Phe  
 530 535 540

Thr Pro Gly Ser Ser Val Ala Ala Ser Trp Leu Trp Gln Leu Ala Lys  
 545 550 555 560

Thr His Val Leu Ala His Asp Thr Gly Tyr His Gln Leu Val Ser His  
 565 570 575

Trp Leu Arg Thr His Cys Cys Val Glu Pro Tyr Val Ile Ala Ala Asn  
 580 585 590

Arg Arg Leu Ser Gln Met His Pro Ile Tyr Arg Leu Leu His Pro His  
 595 600 605

Phe Arg Phe Thr Met Glu Ile Asn Ala Gln Ala Arg Gly Met Leu Ile  
 610 615 620

Asn Ala Asn Gly Ile Ile Glu Ser Ala Phe Ala Pro Gly Lys Leu Cys  
 625 630 635 640

Met Glu Leu Ser Ser Ala Val Tyr Asp Lys Phe Trp Arg Phe Asp Met  
 645 650 655

- 7 -

Glu Ala Leu Pro Ala Asp Leu Ile Arg Arg Gly Met Ala Ile Glu Cys  
                   660                                  665                                  670

Glu Asp Gly Lys Leu Glu Leu Thr Ile Glu Asp Tyr Pro Tyr Ala Asn  
                   675                                  680                                  685

Asp Gly Leu Leu Ile Trp Asp Ser Ile Lys Glu Trp Val Ser Asp Tyr  
                   690                                  695                                  700

Val Asn His Tyr Tyr Gln Leu Ala Ser Asp Ile His Met Asp Lys Glu  
                   705                                  710                                  715                                  720

Leu Gln Gly Trp Trp Asn Glu Val Arg Thr Lys Gly His Pro Asp Lys  
                                   725                                  730                                  735

Glu Glu Gly Trp Pro Glu Leu Asn Cys His Gly Ser Leu Val Glu Val  
                                   740                                  745                                  750

Leu Thr Thr Ile Ile Trp Val Ala Ser Gly His His Ala Ala Val Asn  
                   755                                  760                                  765

Phe Gly Gln Tyr Pro Tyr Ala Gly Tyr Phe Pro Asn Arg Pro Thr Ile  
                   770                                  775                                  780

Ala Arg Arg Asn Met Pro Thr Glu Gly Gln Ala Cys Ser His Asp Gly  
                   785                                  790                                  795                                  800

Met Gln Pro Thr Phe Val Glu Asp Pro Val Arg Val Leu Leu Asp Thr  
                                   805                                  810                                  815

Phe Pro Ser Gln Tyr Gln Thr Thr Leu Val Leu Pro Val Leu Asn Leu  
                                   820                                  825                                  830

Leu Ser Ser His Ser Pro Gly Glu Glu Tyr Met Gly Thr His Ala Glu  
                   835                                  840                                  845

Ser Ala Trp Met Ala Asp Arg Glu Val Arg Ala Ala Phe Gly Arg Phe  
                   850                                  855                                  860

Asn Glu Arg Met Met Ser Ile Ala Glu Met Ile Asp Cys Arg Asn Lys  
                   865                                  870                                  875                                  880

Asp Pro Glu Arg Lys Asn Arg Gln Gly Pro Gly Val Val Pro Tyr Val  
                                   885                                  890                                  895

Leu Leu Lys Pro Ser Tyr Gly Asp Pro Lys Asp Met Thr Ser Val Met  
                                   900                                  905                                  910

Glu Met Gly Ile Pro Asn Ser Ile Ser Ile  
                   915                                  920

&lt;210&gt; 8

&lt;211&gt; 17

- 8 -

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
oligonucleotide

<400> 8  
caygcngtna anttygg

17

<210> 9  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
oligonucleotide

<400> 9  
aatgcttttt tttttttttt v

21

<210> 10  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
oligonucleotide

<400> 10  
gtcagcatgc tcacggccac

20

<210> 11  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:  
oligonucleotide

<400> 11  
cattgacgac ctccgacaag

20

<210> 12  
<211> 37  
<212> DNA  
<213> Artificial Sequence

- 9 -

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 12

caagaaagct gggttgacaa attaagttgt cagtgtg

37

&lt;210&gt; 13

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 13

caaaaaagca ggcttgtaac atcctactcc tattgtg

37

&lt;210&gt; 14

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 14

caaaaaagca ggctccccgt ctttatctac tc

32

&lt;210&gt; 15

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 15

ggggacaagt ttgtacaaaa aagcaggct

29

&lt;210&gt; 16

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 16

ggggaccact ttgtacaaga aagctgggt

29

&lt;210&gt; 17

&lt;211&gt; 4569

&lt;212&gt; DNA

<213> *Oryza sativa*

&lt;400&gt; 17

```
ctgcaggccc agcaaacacg cgtcacacgt ccccgctctt atctactcgc gcccatcctc 60
tacttatcta ctgggcctct cctcctccac tactcctcgc ctggtcctcc cctctcctcg 120
atccccctct ccccttcccg ggctctcccg tgggtggttg cgcgcggcgg caggcggagg 180
cgggcggtgg tgcgcggcgg cgtgacataa gcgcggggct gatgggaggg cgggcggcgg 240
gtggagggga ggatggtggc gggcggatgg gtgggcgaca acgggtgggc ggcagcggga 300
ggcgggtgcc tccttctg cggaggggag gcgggagggc gcagcggcag cggcggattt 360
ttttgttctt ttttttttag tatgttgat tcattggatc tgagatgtat aatttgtgat 420
gtattttttt ttagaatttg tgatgtgaat ctatgatttg tgatgttact ttgatttggg 480
gatatgtagg gagcaactcg atttgaaaaa atcaaatgc aagcagcaaa gaaacaatga 540
caaaaaaaga agaaaaatgg tgaccaaccg aaactgccgc aaccatcttt taatcccggt 600
tggtaacacc aaccagaact gaagatggac atctttaatt tagtctcgga ttcacgggtc 660
tggtttacaa ccgggacta aagggggttg cgaaccagga ctaaagaagg gttttccagc 720
agttacagaa attaaaagta tatagtattt cgtatatatta ctattttgcc actatgtttg 780
tcggagtgtg tacatagata gggggccatt cgatgggacc gctttctatg agttcggggg 840
tgtcgcacgt gagggagcgc tcgggttccc attgaggacg gaaacaaatg aacaaacaat 900
aattatatta gtttgagaca tccggatgtg taacatccta ctctattgt gtcatgtgtt 960
ttctctcttg ctagtgaatg atcttatatg tcagacttag agtcgtagtt catctctttt 1020
ggatttccga agaacttcac tgtgtatggc tataatgttct aataactctg tatgttcgaa 1080
cctcctttct cttgggtcatg gctctcccat tatactatc tagggacacc acatgatcag 1140
aatattaggg agatgcta atagtcata gtgccggatt atgtaaggaa aatcttgtca 1200
gcactgctcc tctctccctt tttgcgaggc ttagcgtaga accacgaaaa aaaaacttgt 1260
attaaagaca ccaagataac tagctgcaaa catcccagac aagccgatca tactcgaaat 1320
caaccacctt ctctatcaca tcatattgaa ttaagggtt atccaatcac attcaaagt 1380
actaactaca gcaaccagga aaatcctcga aagaattaga agcatcaatg aagaagcatt 1440
gaacatcact tctataatcc ctccaggaaa ccatcagaag taaaacattc tctggccccc 1500
attggttgaa tttttttcta ataagccaaa acggcttatt agagaataaa aataaatgtg 1560
taggtaaaaa ttttatatat gtgttttttt taacttaaaa gccaatactg aaaaaaacta 1620
cgttgaaaat atctcagaat caatctcaaa attaagtttg aaaattcaaa atttggctta 1680
ttctttgggt tattgggcca tctgatggga gcctctatat ttggtagcac ataactaaat 1740
aaataaaaagg gttcatattc ctaattgtcg aaagtctctc aaagtgacat aacacactag 1800
tatttacact acggacaacc ctctagtgtt agactttaga gttacatgtt gccagtttc 1860
gaggtaaagt agactagctc aaatgtgtct caaacagtcg ttcccgggaa ataagtcaca 1920
agtgttccca cgtgtaggca tgtttcacgc ttagatcgat cgagtttcgt ttccatctgt 1980
acgtacgttg taccaccca acccgctgat cgatatgatc gatcgtcccc tataaattgg 2040
acatcctgga catgctcttt ctccagacat ctccaccctt cctatcttga tatataactt 2100
cgacacactg acaacttaat ttgtcagaat gctcacggcc acgcagactc tggcgccggc 2160
agtgtctccc cggagccatg ggcgcccttc ttcttctcc agccagccgc gccgcaccgc 2220
cgccgcccgc tcgagagtaa gctgcacccg cgtcggcgcc ttgtcggagg tcgtcaatgg 2280
cgaactcgtc gtcggcgacc aagaacagac gaccgacgac ctctttacgc ggcacaagaa 2340
tgtcgtcgcc gactacacgc tgagcgccac ggtgacgggt agcttgaagc aggacgattc 2400
cactccccag aaggtggcgg acatggttaa tcgagactgg cttttccttg atttcttcag 2460
ctcgcatata ggtgatgaca agctccatag ctactctatc gatcgtaccg gccgcctctt 2520
tcgcgccata tataatcttg aacatgggca ctgattccca ttcgaaaaaa tatcttaatt 2580
tgagaggatg ctttctactg gtagctagca ttctactact agaattatcg tattaagaat 2640
```

```

gatcgcagcc gttcgaattta ctagctagct cgaatgatag catatattgt aagtgtaaat 2700
tttgatacat ttgatatat aacttcgaca ttaggaaaaa aactacatta aaaatttttg 2760
tacctaataa actataaaat ttctactatt atcagtaaaa atattttttg cttaccatta 2820
gtagattaga ttgtatataa gatatactac ctctgttttt taatagatga cgccgttgac 2880
tttttctcac atgttttgacc attcgtctta ttaaaaaaatt atataattat aattttagttt 2940
gttatgaatt gttttatcac tcatagtact ttaagtgtga tttatatctt atacattttac 3000
ataaaaatttt tgaataagac gaatgggtcaa acatgggaga aaaagtcaac ggcgatcatc 3060
attaaaaaac ggagagagta tatgattcat cggaattaaa aaatagacgg tataacacat 3120
tgtaaaacac ctattttcgt cactctacag aggggatgca cacggagcct cagctcgcca 3180
ggtactcgca catggatggc aaaggctcct tcatatacga ggccagcttc agcatcccgt 3240
cctcgttgga cgccgtcggc gccgtgcagg tcgtgaaccg ctacagcagc gaggtgtaca 3300
tctcggacat cgacgtccac ctctgcggcg gccgccatca gtggaccgac atcactttcc 3360
actgcaactc ttggatcgac tacaaccca acgaccagcg cttcttcttc cctctcaagg 3420
tcaggggttc acgttattgc cgataccgat attgttttat tcgtgaggat ggtaaaatta 3480
gctatcggcc aaaaattcct tttttttttg aattagctca tatgtaaaat aaaattaaaa 3540
cctgcaaacc atcgaaaagg gtggaggatt atatttgctt aattagggca gtcccaaccc 3600
ataacactag acatggttta tataacacta gacatcatca agaaactagt actacacact 3660
actcttccaa tgcaaatacc actattccat acttcagtta aatgctactt atctcacatg 3720
atgtcttaga tgttggtgtag aaaccatgtc tcatgcaaga catggtttcc ttctctttct 3780
ttattttatc acttgtcaca ttataatttt gtcctagggtg gcaacttatt taatgctatg 3840
gacaccattc tagttattgg gttgggaatg cccttaagggt tgtgtttgat tgggtggctg 3900
aagctgcatt ggagttgggg ataaggcagc catctctcac tagattggat gagtggatga 3960
ggagaacacc atgtgggtga tgttgctgat tggctgtgat aggtttaact tgatgttaac 4020
aatctcaaac tgattttttt ttctaaactg attatcctat ttataatccg attatatgat 4080
tatatttata ttatagttaa tcttcaaac aaaatttcac atggttatac gaatgggtgc 4140
cactgactta ttgcgtccat gctatatcct ccagcctatg gatatatcaa acaaaacata 4200
gatgggttat cccatcccca tccacttatg accgtgaact aaacgcgccc taatctccat 4260
ttccgctcga tcgatcgttg ttcagctgca aatcattcac actgcttggt atttgcttgc 4320
atgcacgata ttgcagtcgt acctcccgtc tcagacgccc aggggctgta agaactctgcg 4380
caaggaagag ctcaggggcca tccgcggcga tggccgcggc gagcgcaagg agtgggagcg 4440
catctacgac tacgacgtct acaacgacct cggcgacccc gacaatgacc cggccactcg 4500
tcggccgggtg ctcggcggcc gccggcgccc ctaccgcgc cgctgccgca cgggccgccc 4560
ccgctgcag 4569

```

&lt;210&gt; 18

&lt;211&gt; 1198

&lt;212&gt; DNA

<213> *Oryza sativa*

&lt;400&gt; 18

```

tgtaacatcc tactcctatt gtgtcatgtg ttttctctct tgctagtga tgaatcttata 60
tgtcagactt agagtcgtag ttcatctctt ttggatttcc gaagaacttc actgtgtatg 120
gctatatgtt ctaataactc tgtatgttcg aacctccttt ctcttggtca tggctctccc 180
attatatcta tctagggaca ccacatgatc agaataattag ggagatgcta attaatgtga 240
tagtgccgga ttatgtaagg aaaatcttgt cagcactgct cctctctccc tttttgcgag 300
gcttagcgta gaaccacgaa aaaaaaactt gtattaaaga caccaagata actagctgca 360
aacatcccag acaagccgat catactcgaa atcaaccacc ttctctatca catcatattg 420
aatttaagg tttatccaatc acattcaaaag ttactaacta cagcaaccag gaaaatcctc 480
gaaagaatta gaagcatcaa tgaagaagca ttgaacatca cttctataat ccctccagga 540
aaccatcaga agtaaaacat tctctggccc ccattgggtg aatttttttc taataagcca 600
aaacggctta ttagagaata aaaataaatg tgtaggtaaa acttttatat atgtgttttt 660
tttaacttaa aagccaatac tgaaaaaac tacgttgaaa atatctcaga atcaatctca 720
aaattaagtt tgaaaattca aaatttggtc tattctttgg cttattggggc catctgatgg 780
gagcctctat atttggtagc acataactaa ataaataaaa gggttcatat tcctaattgt 840

```

- 12 -

```

cgaaagtctc tcaaagtgc ataacacact agtattttaca ctacggacaa ccctcctagt 900
ttagacttta gagttacatg ttgccagtt tcgaggtaaa gtagactagc tcaaagtgtg 960
ctcaaacagt cgttcccgga acataagtca caagtgttcc cactgttagg catgtttcac 1020
gcttagatcg atcgagtttc gtttccatct gtacgtacgt tgtaccaccc caaccgcgcg 1080
atcgatatga tcgatcgtcc cctataaatt ggacatcctg gacatgctct ttcttcagac 1140
atctccaccc ttcctatctt gatataatac ttgcacacac tgacaactta atttgtca 1198

```

&lt;210&gt; 19

&lt;211&gt; 2096

&lt;212&gt; DNA

<213> *Oryza sativa*

&lt;400&gt; 19

```

ccccgtcttt atctactcgc gcccatcctc tacttatcta ctcggcctct cctcctccac 60
tactcctcgc ctggtcctcc cctctcctcg atccccctct ccccttcccg ggctctcccg 120
tggtgggttg cgcgcggcgg caggcggagg cgggcgggtg tgcgcggcgg cgtgacataa 180
gcggcgggct gatgggaggg cgggcggcgg gtggagggga ggatggtggc gggcggatgg 240
gtgggcgaca acgggtgggc ggcagcggga ggcggtgccc tcccttctgc cggaggggag 300
gcgggaggcg gcagcggcag cggcggattt ttttgttctt ttttttttag tatttgtgat 360
tcattggatc tgagatgtat aatttgtgat gtattttttt ttagaatttg tgatgtgaat 420
ctatgatttg tgatgttact ttgatttggg gatatgtagg gagcaactcg atttggaaaa 480
atcaaaatgc aagcagcaaa gaaacaatga caaaaaaaga agaaaaatgg tgaccaaccg 540
aaactgccgc aaccatcttt taatcccggg tggtaacacc aaccagaact gaagatggac 600
atctttaatt tagtctcgga ttcacggtcc tggtttacia cccgggacta aagggggttg 660
cgaaccagga ctaaaagaagg gttttccagc agttacagaa attaaaagta tatagtattt 720
gctatattta ctattttgcc actatgtttg tcggagtgtg tacatagata gggggccatt 780
cgatgggacc gctttctatg agttcggggg tgtcgcacgt gagggagcgc tcgggttccc 840
attgaggacg gaaacaaatg aacaaacaat aattatatta gtttgagaca tccggatgtg 900
taacatccta ctctattgtg gtcatgtgtt ttctctcttg ctagtgaatg atcttatatg 960
tcagacttag agtcgtagtt catctctttt ggatttccga agaacttcac tgtgtatggc 1020
tatatgttct aataactctg tatgttcgaa cctcctttct cttgggtcatg gctctcccat 1080
tatatctatc tagggacacc acatgatcag aatattaggg agatgctaata agatgcata 1140
gtgccggatt atgtaaggaa aatcttgtca gcaactgctc tctctccctt ttgcgaggc 1200
ttagcgtaga accacgaaaa aaaaacttgt attaaagaca ccaagataac tagctgcaaa 1260
catcccagac aagccgatca tactcgaaat caaccacctt ctctatcaca tcatattgaa 1320
ttaaaggttt attccaatcac attcaaagtt actaactaca gcaaccagga aaatcctcga 1380
aagaattaga agcatcaatg aagaagcatt gaacatcact tctataatcc ctccaggaaa 1440
ccatcagaag taaaacattc tctggcccc attggttgaa tttttttcta ataagccaaa 1500
acggcttatt agagaataaa aataaatgtg taggtaaaac ttttatatat gtgttttttt 1560
taacttaaaa gccaatactg aaaaaaacta cgttgaaaat atctcagaat caatctcaaa 1620
attaagtttg aaaattcaaa atttggctta ttctttggct tattgggcca tctgatggga 1680
gcctctatat ttggtagcac ataactaaat aaataaaaagg gttcatattc ctaattgtcg 1740
aaagtctctc aaagtgcac aacacactag tatttacact acggacaacc ctctagttt 1800
agactttaga gttacatgtt gccagtttc gaggtaaaag agactagctc aaatgtgtct 1860
caaacagtcg ttcccggaac ataagtcaca agtggtccca cgtgtaggca tgtttcacgc 1920
ttagatcgat cgagtttcgt ttccatctgt acgtacgttg taccacccca acccgtcgat 1980
cgatatgac gatcgtcccc tataaattgg acatcctgga catgctcttt ctccagacat 2040
ctccaccctt cctatcttga tatataactt cgacacactg acaacttaat ttgtca 2096

```

&lt;210&gt; 20

&lt;211&gt; 13274

&lt;212&gt; DNA

&lt;213&gt; artificial

&lt;400&gt; 20

ccaattcccg	atctagtaac	atagatgaca	ccgcgcgcga	taatttatcc	tagtttgccg	60
gctatatttt	gttttctatc	gcgtattaaa	tgtataattg	cgggactcta	atcataaaaa	120
cccatctcat	aaataacgtc	atgcattaca	tgttaattat	tacatgctta	acgtaattca	180
acagaaatta	tatgataatc	atcgcaagac	cggcaacagg	attcaatctt	aagaaacttt	240
attgccaaat	gtttgaacga	tcggggaaat	tcggggatct	aattcccgag	gctgtagccg	300
acgatgggtc	gccaggagag	ttgttgattc	attgtttgcc	tccctgctgc	ggtttttcac	360
cgaagtcat	gccagtcag	cgtttttgca	gcagaaaagc	cggcgacttc	ggtttgcggt	420
cgcgagtga	gatccctttc	ttgttaccgc	caacgcgcaa	tatgccttgc	gaggtcgcaa	480
aatcggcgaa	attccatacc	tgttcaccga	cgacggcgct	gacgcgatca	aagacgcggt	540
gatacatatc	cagccatgca	cactgatact	cttcactcca	catgtcgggt	tacattgagt	600
gcagcccggc	taacgtatcc	acgcggtatt	cgggtgatgat	aatcggctga	tgcagtttct	660
cctgccaggc	cagaagttct	ttttccagta	ccttctctgc	cgtttccaaa	tcgccgcttt	720
ggacatacca	tccgtaataa	cggttcaggc	acagcacatc	aaagagatcg	ctgatgggat	780
cgggtgtgagc	gtcgcagaac	attacattga	cgcagggtgat	cggacgcgtc	gggtcgagtt	840
tacgcgttgc	ttccgccagt	ggcgcgaaat	attcccgtgc	accttgcgga	cgggtatccg	900
gttcggttgc	aatactccac	atcaccacgc	ttgggtgggt	ttgtcacgc	gctatcagct	960
ctttaatcgc	ctgtaagtgc	gcttgctgag	tttcuccgtt	gactgcctct	tcgctgtaca	1020
gttctttcgg	cttgttgccc	gcttcgaaac	caatgcctaa	agagagggtta	aagccgacag	1080
cagcagtttc	atcaatcacc	acgatgccat	gttcatctgc	ccagtcgagc	atctcttcag	1140
cgtaagggtta	atgcgaggta	cggtaggagt	tggccccaat	ccagtcatt	aatgcgtggt	1200
cgtgcaccat	cagcacgtta	tcgaatcctt	tgccacgcaa	gtccgcatct	tcatgacgac	1260
caaagccagt	aaagtagaac	ggtttgtggt	taatcaggaa	ctgttcgccc	ttactgcca	1320
ctgaccggat	gccgacgcga	agcgggtaga	tatcacactc	tgtctggctt	ttggctgtga	1380
cgcacagttc	atagagataa	ccttcacccg	gttgccagag	gtgcggatc	accacttgca	1440
aagtcgccgt	agtgccttgt	ccagttgcaa	ccacctgttg	atccgcatca	cgcagttcaa	1500
cgctgacatc	accattggcc	accacctgcc	agtcaacaga	cgcgtgggtta	cagtcttgccg	1560
cgacatgcgt	caccacgggt	atatcgcca	cccagggtgt	cggcgtgggt	tagagcatta	1620
cgctgcgatg	gattccggca	tagttaaaga	aatcatggaa	gtaagactgc	ttttcttgc	1680
cgttttcgtc	ggtaatcacc	attcccggcg	ggatagctctg	ccagttcagt	tcgttgttca	1740
cacaaacggt	gatacctgca	catcaacaaa	ttttggtcat	atattagaaa	agttataaat	1800
taaaatatac	acacttataa	actacagaaa	agcaattgct	atatactaca	ttcttttatt	1860
ttgaaaaaaa	tatttgaaat	attatatatt	tactaattaa	tgataattat	tatatata	1920
tcaaaggtag	aagcagaaac	ttacgtacac	ttttcccgcc	aataacatac	ggcgtgacat	1980
cggcttcaaa	ttggcgtag	ccgcctgat	gctccatcac	ttctgatta	ttgaccaca	2040
ctttgccgta	atgagtgacc	gcatcgaaac	gcagcacgat	acgtggcct	gccaacctt	2100
tcggtataaa	gacttcgcgc	tgataccaga	cgttgccgcg	ataattacga	atatctgcat	2160
cggcgaactg	atcgttaaaa	ctgcctggca	cagcaattgc	cgggttttct	tgtaacgcgc	2220
tttcccacca	acgtgatca	attccacagt	tttcgcgatc	cagactgaat	gcccacaggc	2280
cgctgagttt	tttgatttca	cgggttgggg	tttctacagg	acggaccatg	gtcgacctcg	2340
aatcaaccac	tttgatacaag	aaagctgaac	gagaaacgta	aatgatata	aatatcaata	2400
tattaaatta	gattttgcat	aaaaaacaga	ctacataata	ctgtaaaaca	caacatatcc	2460
agtcactatg	gtcgacctgc	agactggctg	tgtataaggg	agcctgacat	ttatattccc	2520
cagaacatca	gggttaatgyc	gtttttgatg	tcattttcgc	ggtggctgag	atcagccact	2580
tcttccccga	taacggagac	cggcacactg	gccatatcgg	tggatcatcat	gcgccagctt	2640
tcatccccga	tatgaccac	cgggttaaagt	tcacgggaga	ctttatctga	cagcagacgt	2700
gcactggcca	gggggatcac	catccgtcgc	cggggcgtgt	caataatata	actctgtaca	2760
tccacaaaca	gacgataacg	gctctctctt	ttataggtgt	aaaccttaaa	ctgcatttca	2820
ccagcccctg	ttctcgtcag	caaaagagcc	gttcatttca	ataaacgggg	cgacctcagc	2880
catcccttcc	tgattttccg	ctttccagcg	ttcggcacgc	agacgacggg	cttcattctg	2940
catggttgtg	cttaccagac	cggagatatt	gacatcatat	atgccttgag	caactgatag	3000
ctgcgctgt	caactgtcac	tgtaatacgc	gtcttcatag	catacctctt	tttgacatac	3060
ttcgggtata	catatcagta	tatattctta	taccgcaaaa	atcagcgccg	aaatacgcac	3120
actgttatct	ggcttttagt	aagccggatc	cagatcttta	cgcgccgcc	tgccactcat	3180



cgcagtactg	ttgtaattca	ttaagcattc	tgccgacatg	gaagccatca	caaacggcat	3240
gatgaacctg	aatcgccagc	ggcatcagca	ccttgtcgcc	ttgctataa	tatttgccca	3300
tggtgaaaac	gggggcgaag	aagttgtcca	tattggccac	gtttaaatca	aaactggtga	3360
aaactaccca	gggattggct	gagacgaaaa	acataattctc	aataaacctt	ttagggaaat	3420
aggccagggt	ttcaccgtaa	cacgccacat	ccttgcgaata	tatgtgtaga	aactgccgga	3480
aatcgtcgtg	gtattcactc	cagagcgatg	aaaacgtttc	agtttgcctc	tggaaaacgg	3540
tgtaacaagg	gtgaacacta	tcccatatca	ccagctcacc	gtctttcatt	gccatacggg	3600
attccggatg	agcattcatc	aggcgggcaa	gaatgtgaat	aaaggccgga	taaaacttgt	3660
gcttattttt	ctttacggtc	tttaaaaagg	ccgtaatatc	cagctgaacg	gtctgggtat	3720
aggtacattg	agcaactgac	tgaaatgcct	caaaatgttc	tttacgatgc	cattgggata	3780
tatcaacggg	ggtatatcca	gtgatttttt	tctccatttt	agcttcctta	gctcctgaaa	3840
atctcgacgg	atcctaactc	aaaatccaca	cattatacga	gccggaagca	taaagtgtaa	3900
agcctggggg	gcctaatacg	gcgcccatag	tgactggata	tgttgtgttt	tacagtatta	3960
tgtagtctgt	tttttatgca	aaatctaatt	taatatattg	atatttatat	cattttacgt	4020
ttctcgttca	gcttttttgt	acaaacttgt	tgattcgagg	ggatcctcta	gagtcgacct	4080
gcaggcatgc	aaagctcggg	accagcttgc	atgcctgcag	tgacgcgtga	cccggtcgtg	4140
cccctctcta	gagataatga	gcattgcatg	tctaagttat	aaaaaattac	cacatatttt	4200
ttttgtcaca	cttgtttgaa	gtgcagttta	tctatcttta	tacatatatt	taaactttac	4260
tctacgaata	atataatcta	tagtactaca	ataaatcag	tgttttagag	aatcatataa	4320
atgaacagtt	agacatggtc	taaaggacaa	ttgagtattt	tgacaacagg	actctacagt	4380
tttatctttt	tagtgtgcat	gtgttctcct	ttttttttgc	aaatagcttc	acctatataa	4440
tacttcaccc	atgtttattag	tacatccatt	taggggttag	ggttaatggg	ttttatagac	4500
taattttttt	agtacatcta	ttttattcta	ttttagcctc	taaattaaga	aaactaaaac	4560
tctatttttag	tttttttatt	taataattta	gatataaaaat	agaataaaaat	aaagtgacta	4620
aaaattaaac	aaataccctt	taagaaatta	aaaaaactaa	ggaaacattt	ttcttgtttc	4680
gagtagataa	tgccagcctg	ttaaacgccc	tcgacgagtc	taacggacac	caaccagcga	4740
accagcagcg	tcgctcggg	ccaagcgaag	cagacggcac	ggcatctctg	tcgctgcctc	4800
tggaaccctc	tcgagagttc	cgctccaccg	ttggacttgc	tccgctgtcg	gcacccagaa	4860
attgctgggc	ggagcggcag	acgtgagccg	gcacggcagg	cggcctcctc	ctcctctcac	4920
ggcacgggca	gctacggggg	attcctttcc	caccgctcct	tcgctttccc	ttcctcgccc	4980
gccgtaataa	atagacaccc	cctccacacc	ctctttcccc	aacctcgtgt	tggtcggagc	5040
gcacacacac	acaaccagat	ctcccccaaa	tccaccgcgc	ggcacctccg	cttcaaggta	5100
cgccgctcgt	cctccccccc	ccccctctc	tacctctctc	agatcggcgt	tccggtccat	5160
ggttaggggc	cggtagttct	acttctgttc	atgtttgtgt	tagatccgtg	tttgtgttag	5220
atccgtgctg	ctagcgttcg	tacacggatg	cgacctgtac	gtcagacacg	ttctgattgc	5280
taacttgcca	gtgtttctct	ttggggaatc	ctgggatggc	tctagccgtt	ccgcagacgg	5340
gatcgatttc	atgatttttt	ttgtttcgtt	gcatagggtt	tggtttgccc	ttttccttta	5400
tttcaatata	tgcggtgcac	ttgtttgtcg	ggcatccttt	tcatgctttt	ttttgtcttg	5460
gttgtgatga	tggtgtctgg	ttgggcgggt	gttctagatc	ggagtagaat	tctgtttcaa	5520
actacctggg	ggattttatta	atgtttggatc	gttatgtgtg	tgccatacat	attcatagtt	5580
acgaattgaa	gatgatggat	ggaaatatcg	atctaggata	ggtatacatg	ttgatgcggg	5640
ttttactgat	gcataacacg	agatgctttt	tgttcgcttg	gttgtgatga	tgtggtgtgg	5700
ttgggcgggc	gttcattcgt	tctagatcgg	agtagaatac	tgtttcaaac	tacctggtgt	5760
atttattaat	tttggaactg	tatgtgtgtg	tcatacatct	tcatagttac	gagtttaaga	5820
tggtatggaaa	tatcgatcta	ggataggtat	acatgttgat	gtgggtttta	ctgatgcata	5880
tacatgatgg	catatgcagc	atctattcat	atgctctaac	cttgagtacc	tatctattat	5940
aataaacaag	tatgttttat	aattattttg	atcttgatat	acttggatga	tggcatatgc	6000
agcagctata	tgtggatttt	tttagccctg	ccttcatacg	ctattttatt	gcttgggtact	6060
gtttcttttg	tcgatgctca	ccctgttgtt	tggtgttact	tctgcaggga	tccccgatca	6120
tgcaaaaact	cattaactca	gtgcaaaaact	atgcctgggg	cagcaaaaacg	gcgttgactg	6180
aactttatgg	tatggaaaat	ccgtccagcc	agccgatggc	cgagctgtgg	atgggcgcac	6240
atccgaaaag	cagttcacga	gtgcagaatg	ccgcccggaga	tatcgtttca	ctgcgtgatg	6300
tgattgagag	tgataaatcg	actctgctcg	gagaggccgt	tgccaaacgc	tttggcgaac	6360
tgcccttctc	gttcaaagta	ttatgcgcag	cacagccact	ctccattcag	gttcattcaa	6420
acaaaacaaa	ttctgaaatc	ggttttgcca	aagaaaatgc	cgcaggtatc	ccgatggatg	6480

ccgccgagcg	taactataaa	gacctaacc	acaagccgga	gctgggtttt	gcgctgacgc	6540
ctttccttgc	gatgaacgcg	tttcgtgaat	tttccgagat	tgtctcccta	ctccagccgg	6600
tcgcaggtgc	acatccggcg	attgctcact	ttttacaaca	gcctgatgcc	gaacgtttaa	6660
gcgaactgtt	cgccagcctg	ttgaatatgc	aggggtgaaga	aaaatcccgc	gcgctggcga	6720
ttttaaaatc	ggccctcgat	agccagcagg	gtgaaccgtg	gcaaaccgatt	cgtttaattt	6780
ctgaatttta	cccggaagac	agcgggtctgt	tctccccgct	attgctgaat	gtgggtgaaat	6840
tgaaccctgg	cgaagcgatg	ttcctgttcg	ctgaaacacc	gcacgcttac	ctgcaaggcg	6900
tggcgctgga	agtgatggca	aactccgata	acgtgctgcg	tgcgggtctg	acgcctaaat	6960
acattgatat	tccggaactg	gttgccaatg	tgaaattcga	agccaaaccg	gctaaccagt	7020
tgttgacca	gccgggtgaa	caaggtgcag	aactggactt	cccgattcca	gtggatgatt	7080
ttgccttctc	gctgcatgac	cttagtgata	aagaaaccac	cattagccag	cagagtgccg	7140
ccattttgtt	ctgcgtcgaa	ggcgatgcaa	cgttgtggaa	aggttctcag	cagttacagc	7200
ttaaaccggg	tgaatcagcg	tttattgccg	ccaacgaatc	accggtgact	gtcaaaggcc	7260
acggccggtt	agcgcgtgtt	tacaacaagc	tgtaagagct	tactgaaaaa	attaacatct	7320
cttgctaagc	tgggagctcg	atccgtcgac	ctgcagatcg	ttcaaacatt	tggcaataaa	7380
gtttcttaag	attgaatcct	gttgccggtc	ttgcgatgat	tatcatataa	tttctgttga	7440
attacgttaa	gcatgtaata	attaacatgt	aatgcatgac	gttatttatg	agatgggttt	7500
ttatgattag	agtcccgcga	ttatacattt	aatacgcgat	agaaaacaaa	atatagccgc	7560
caaactagga	taaattatcg	cgccgggtgt	catctatgtt	actagatctg	ctagccctgc	7620
aggaaattta	ccggtgcccg	ggcgcccgagc	atggccgtat	ccgcaatgtg	ttattaagtt	7680
gtctaagcgt	caatttggtt	acaccacaat	atatcctgcc	accagccagc	caacagctcc	7740
ccgaccggca	gctcggcaca	aaatcaccac	tcgatacagg	cagcccatca	gaattaattc	7800
tcatgtttga	cagcttatca	tcgactgcac	ggtgaccaa	tgcttctggc	gtcaggcagc	7860
catcggaagc	tgtggtatgg	ctgtgcaggt	cgtaaatacac	tgcataattc	gtgtcgctca	7920
aggcgactc	ccgttctgga	taatgttttt	tgccgcgaca	tcataacggg	tctggcaaat	7980
attctgaaat	gagctgttga	caattaatca	tccggctcgt	ataatgtgtg	gaattgtgag	8040
cggataacaa	tttcacacag	gaaacagacc	atgaggggaag	cgttgatcgc	cgaagtatcg	8100
actcaactat	cagaggtagt	tggcgtcatc	gagcgccatc	tcgaaccgac	gttgctggcc	8160
gtacatttgt	acggctccgc	agtggatggc	ggcctgaagc	cacacagtga	tattgatttg	8220
ctggttacgg	tgaccgtaag	gcttgatgaa	acaacgcggc	gagctttgat	caacgacctt	8280
ttggaaactt	cggcttcccc	tggagagagc	gagattctcc	gcgctgtaga	agtcaccatt	8340
gttggtgcacg	acgacatcat	tccgtggcgt	tatccagcta	agcgcgaact	gcaatttggg	8400
gaatggcagc	gcaatgacat	tcttgacaggt	atcttcgagc	cagccacgat	cgacattgat	8460
ctggctatct	tgctgacaaa	agcaagagaa	catagcggtg	ccttggtagg	tccagcggcg	8520
gaggaaactct	ttgatccggt	tcctgaacag	gatctatttg	aggcgctaaa	tgaaacctta	8580
acgctatgga	actcgccgcc	cgactgggct	ggcgatgagc	gaaatgtagt	gcttacgttg	8640
tcccgcattt	ggtacagcgc	agtaaccggc	aaaactcgcc	cgaaggatgt	cgctgcgcag	8700
tgggcaatgg	agcgctgccc	ggcccagtat	cagcccgctca	tacttgaagc	taggcaggct	8760
tatcttggag	aagaagatcg	cttggcctcg	cgcgagatc	agttggaaga	atttgttcac	8820
tacgtgaaag	gcgagatcac	caaagtagtc	ggcaataaaa	gctctagtgg	atctccgtac	8880
ccccggggga	tctggctcgc	ggcgacgca	cgacgcggg	gcgagaccat	aggcgatctc	8940
ctaaatcaat	agtagctgta	acctcgaagc	gtttcacttg	taacaacgat	tgagaatttt	9000
tgtcataaaa	ttgaaatact	tggttcgcac	ttttgtcatc	cgcggtcagc	cgcaattctg	9060
acgaactgcc	catttagctg	gagatgattg	tacatccttc	acgtgaaaat	ttctcaagcg	9120
ctgtgaacaa	gggttcagat	tttagattga	aagggtgagcc	gttgaaacac	gttcttcttg	9180
tcgatgacga	cgtcgctatg	cggcattcta	ttattgaata	ccttacgac	cacgccttca	9240
aagtgaccgc	ggtagccgac	agcaccaggt	tcacaagagt	actctcttcc	gcgacggctg	9300
atgtcgtggg	tgttgatcta	aatttaggtc	gtgaagatgg	gctcgagatc	gttcgtaatc	9360
tggcggcaaa	gtctgatatt	ccaatcataa	ttatcagtg	cgaccgcctt	gaggagacgg	9420
ataaagtgtg	tgactcgag	ctaggagcaa	gtgattttat	cgctaagccg	ttcagtatca	9480
gagagtttct	agcacgcatt	cggtttgcct	tgcgcgtgcg	ccccaacgtt	gtccgctcca	9540
aagaccgacg	gtctttttgt	tttactgact	ggacacttaa	tctcaggcaa	cgctcgttga	9600
tgccgaagc	tggcggtgag	gtgaaactta	cggcaggtga	gttcaatctt	ctcctcgctg	9660
ttttagagaa	accccgcgac	gttctatcgc	gcgagcaact	tctcattggc	agtcgagtac	9720
cgcacgagga	ggtttatgac	aggagtatat	atgttctcat	tttgaggctg	cgccgcaaac	9780

ttgaggcaga	tccgtcaagc	cctcaactga	taaaaacagc	aagaggtgcc	ggttatttct	9840
ttgacgcgga	cgtgcaggtt	tgcacgggg	ggacgatggc	agcctgagcc	aattcccaga	9900
tcccagagga	atcggcggtga	goggtcgcaa	accatccggc	ccggtacaaa	tccggcgggc	9960
gtgggtgat	gacctggtgg	agaagttgaa	ggccgcgcag	gccgccagc	ggcaacgcat	10020
cgaggcagaa	gcacgccccg	gtgaatcgtg	gcaagcgggc	gctgatcgaa	tccgcaaaga	10080
atcccggcaa	ccgccggcag	cgggtgcgcc	gtcgattagg	aagccgcca	agggcgacga	10140
gcaaccagat	tttttcgttc	cgatgctcta	tgacgtgggc	acccgcgata	gtcgcagcat	10200
catggacgtg	gccgttttcc	gtctgtcgaa	gcgtgacoga	cgagctggcg	aggtgatccg	10260
ctacgagctt	ccagacgggc	acgtagaggt	ttccgcaggg	ccggccggca	tggccagtgt	10320
gtgggattac	gacctggtac	tgatggcggt	ttcccatcta	accgaatcca	tgaaccgata	10380
ccgggaaggg	aagggagaca	agcccgccg	cgtgttccgt	ccacacgttg	cggacgtact	10440
caagttctgc	cggcgagccg	atggcgga	gcagaaagac	gacctggtag	aaacctgcat	10500
tccgttaaac	accacgcacg	ttgccatgca	gcgtacgaag	aaggccaaga	acggccgcct	10560
ggtgacggta	tccgaggggtg	aagccttgat	tagccgctac	aagatcgtaa	agagcgaaac	10620
cgggcgcccg	gagtacatcg	agatcgagct	agctgattgg	atgtaccgcg	agatcacaga	10680
aggcaagaac	ccggacgtgc	tgacggttca	ccccgattac	tttttgatcg	atcccggcat	10740
cggcggtttt	ctctaccgcc	tggcacgccg	cgccgcaggc	aaggcagaag	ccagatgggt	10800
gttcaagacg	atctacgaac	gcagtggcag	cgccggagag	ttcaagaagt	tctgtttcac	10860
cgtgcgaag	ctgatcgggt	caaatacct	gccggagtac	gatttgaagg	aggaggcggg	10920
gcaggctggc	ccgatcctag	tcatgcgcta	ccgcaacctg	atcgagggcg	aagcatccgc	10980
cgttctctaa	tgtacggagc	agatgctagg	gcaaattgcc	ctagcagggg	aaaaaggctc	11040
aaaaggctct	tttcctgtgg	atagcacgta	cattgggaac	ccaaagccgt	acattgggaa	11100
ccggaacccg	taccattggga	acccaaagcc	gtacattggg	aaccggtcac	acatgtaagt	11160
gactgatata	aaagagaaaa	aaggcgattt	ttccgcctaa	aactctttta	aacttattaa	11220
aactcttaaa	acccgcctgg	cctgtgcata	actgtctggc	cagcgcacag	ccgaagagct	11280
gcaaaaagcg	cctacccttc	ggtcgctgcg	ctccctacgc	cccgccgctt	cgctcgcc	11340
tatcgccggc	gctggccgct	caaaaatggc	tggcctacgg	ccaggcaatc	taccagggcg	11400
cggacaagcc	gcgccgtcgc	cactcgaccg	ccggcgctga	ggtctgcctc	gtgaagaagg	11460
tggtgctgac	tcataccagg	cctgaatcgc	cccatcatcc	agccagaaag	tgaggagacc	11520
acggttgatg	agagctttgt	tgtaggtgga	ccagttgggt	attttgaact	tttgctttgc	11580
cacggaacgg	tctgcgttgt	cggaagatg	cgtgatctga	tccttcaact	cagcaaaagt	11640
tcgatttatt	caacaaagcc	gccgtcccgt	caagtcagcg	taatgctctg	ccagtgttac	11700
aaccaattaa	ccaattctga	ttagaaaaac	tcatcgagca	tcaaatgaaa	ctgcaattta	11760
ttcatatcag	gattatcaat	accatatttt	tgaaaaagcc	gtttctgtaa	tgaaggagaa	11820
aactcacoga	ggcagttcca	taggatggca	agatcctggg	atcgggtctgc	gattccgact	11880
cgccaacat	caatacaacc	tattaatttc	ccctcgtcaa	aaataagggt	atcaagtga	11940
aatcaccat	gagtgacgac	tgaatccggt	gagaattggc	aaagctctgc	attaatgaat	12000
cggccaacgc	gcggggagag	gcggtttgcg	tattggggcg	tcttcgctt	cctcgctcac	12060
tgactcgctg	cgctcggtcg	ttcggtcgcg	gcgagcggtg	tcagctcact	caaaggcggt	12120
aatacggtta	tccacagaat	caggggataa	cgcaggaaag	aacatgtgag	caaaaggcca	12180
gcaaaaggcc	aggaaccgta	aaaaggccgc	ggtgctggcg	tttttcata	ggctccgccc	12240
ccctgacgag	catcacaaaa	atcgacgctc	aagtcagagg	tggcgaaacc	cgacaggact	12300
ataaagatac	caggcggttc	cccctggaag	ctccctcggt	cgctctcctg	ttccgacct	12360
gccgcttacc	ggatacctgt	ccgcctttct	cccttcggga	agcgtggcgc	tttctcatag	12420
ctcacgctgt	aggtatctca	gttcggtgta	ggtcggtcgc	tccaagctgg	gctgtgtgca	12480
cgaaccccc	gttcagcccg	accgctgcgc	cttatccggt	aactatcgtc	ttgagtccaa	12540
cccggtgaaga	cacgacttat	cgccactggc	agcagccact	ggtaacagga	ttagcagagc	12600
gaggtatgta	ggcggtgcta	cagagttctt	gaagtgggtg	cctaactacg	gctacactag	12660
aagaacagta	tttggatatct	gcgctctgct	gaagccagtt	accttcggaa	aaagagttgg	12720
tagctcttga	tccggcaaac	aaaccaccgc	tggtagcggg	ggtttttttg	tttgcaagca	12780
gcagattacg	cgcagaaaaa	aaggatctca	agaagatcct	ttgatctttt	ctacggggtc	12840
tgaogctcag	tggaaacgaa	actcacgtta	agggattttg	gtcatgagat	tatcaaaaag	12900
gatcttcacc	tagatccttt	tgatccggaa	ttaattcctg	tggttggcat	gcacatacaa	12960
atggacgaac	ggataaacct	tttcacgccc	ttttaaatat	ccgattatct	taataaacgc	13020
tcttttctct	taggtttacc	cgccaatata	tccgtgcaaa	cactgatagt	ttaaactgaa	13080

- 17 -

ggcgggaaac	gacaatctga	tcatgagcgg	agaattaagg	gagtcacgtt	atgacccccg	13140
ccgatgacgc	gggacaagcc	gttttacgtt	tggaactgac	agaaccgcaa	cgctgcagga	13200
attggccgca	gcgccattt	aaatcaattg	ggcgcgcgca	attcgagctc	ggtacaagct	13260
tggcgcgcg	gtac					13274

&lt;210&gt; 21

&lt;211&gt; 2001

&lt;212&gt; DNA

&lt;213&gt; artificial

&lt;400&gt; 21

atggtccgtc	ctgtagaaac	cccaaccggt	gaaatcaaaa	aactcgacgg	cctgtgggca	60
ttcagctctg	atcgcgaaaa	ctgtggaatt	gatcagcgtt	ggtgggaaag	cgcgttacaa	120
gaaagccggg	caattgctgt	gccaggcagt	tttaacgac	agttcgccga	tgcagatatt	180
cgtaattatg	cgggcaacgt	ctggtatcag	cggaagtct	ttataccgaa	aggttgggca	240
ggccagcgtg	tctgtctgct	tttcgatgct	gtcactcatt	acggcaaatg	gtgggtcaat	300
aatcagggaag	tgatggagca	tcaggggcgg	tatacgccat	ttgaagccga	tgtcacgccg	360
tatgttattg	ccgggaaaag	tgtacgtaag	tttctgcttc	tacctttgat	atataataa	420
taattatcat	taattagtag	taataataa	tttcaaata	ttttttcaa	ataaaagaat	480
gtagtatata	gcaattgctt	ttctgtagtt	tataagtgtg	tatattttaa	tttataactt	540
ttctaataa	tgacaaaaat	ttgttgatgt	gcaggatata	ccgtttgtgt	gaacaacgaa	600
ctgaactggc	agactatccc	gccgggaatg	gtgattaccg	acgaaaacgg	caagaaaaag	660
cagctcttact	tccatgattt	ctttaactat	gccggaatcc	atcgacgctt	aatgctctac	720
accacgcccga	acacctgggt	ggacgatata	accgtggtga	cgcatgtcgc	gcaagactgt	780
aaccacgctg	ctgttgactg	gcagggtggt	gccaatgggt	atgtcagcgt	tgaactgcgt	840
gatgcggatc	aacagggtgt	tgcaactgga	caaggcacta	gcgggacttt	gcaagtgggt	900
aatccgcacc	tctggcaacc	gggtgaaggt	tatctctatg	aactgtgcgt	cacagccaaa	960
agccagacag	agtgtgatat	ctaccgcgtt	cgctgcggca	tccggtcagt	ggcagtgaag	1020
ggcgaacagt	tcttgattaa	ccacaaaccg	ttctacttta	ctggcttttg	tctgcatgaa	1080
gatgcggact	tgcgtggcaa	aggattcgat	aacgtgctga	tggtgcacga	ccacgcatta	1140
atggactgga	ttggggccaa	ctcctaccgt	acctcgcat	accttacgc	tgaagagatg	1200
ctcgactggg	cagatgaaca	tggcatcgtg	gtgattgatg	aaactgctgc	tgtcggcttt	1260
aacctctctt	taggcattgg	tttcgaagcg	ggcaacaagc	cgaaagaact	gtacagcgaa	1320
gaggcagtc	acggggaaac	tcagcaagcg	cacttacagg	cgattaaaga	gctgatagcg	1380
cgtgacaaaa	accacccaag	cgtggtgatg	tggagtattg	ccaacgaacc	ggatacccg	1440
ccgcaagggt	cacgggaata	tttcgcgcca	ctggcggaag	caacgcgtaa	actcgaccg	1500
acgcgtccga	tcacctgcgt	caatgtaatg	ttctgcgacg	ctcacaccga	taccatcagc	1560
gatctctttg	atgtgctgtg	cctgaaccgt	tattacggat	ggatgtcca	aagcggcgat	1620
ttggaacgg	cagagaaggt	actggaaaaa	gaacttctgg	cctggcagga	gaaactgcat	1680
cagccgatta	tcatcaccga	atacggcgtg	gatacgttag	ccgggctgca	ctcaatgtac	1740
accgacatgt	ggagtgaaga	gtatcagtg	gcattggctg	atatgtatca	ccgcgtcttt	1800
gatcgcgtca	gcgccgtcgt	cggtgaacag	gtatggaatt	tgcgccattt	tgcgacctcg	1860
caaggcatat	tgcgcgttgg	cggttaacaag	aaagggatct	tactcgcga	ccgcaaaccg	1920
aagtcggcgg	cttttctgct	gcaaaaacgc	tggactggca	tgaacttcgg	tgaaaaaccg	1980
cagcaggag	gcaaacaatg	a				2001

&lt;210&gt; 22

&lt;211&gt; 12817

&lt;212&gt; DNA

&lt;213&gt; artificial

&lt;400&gt; 22

ttttacaaa	cttgttgatt	cgaggggatc	ctctagagtc	gacctgcagg	catgcaaacg	60
-----------	------------	------------	------------	------------	------------	----

tccgtaccag	cttgcattgcc	tgcagtgcag	cgtgaccg	tcgtgcccct	ctctagagat	120
aatgagcatt	gcatgtctaa	gttataaaaa	attaccacat	atTTTTTTTg	tcacacttgt	180
ttgaagtga	gtttatctat	ctttatacat	atatttaaac	tttactctac	gaataatata	240
atctatagta	ctacaataat	atcagtgttt	tagagaatca	tataaatgaa	cagtttagaca	300
tggctctaaag	gacaattgag	tattttgaca	acaggactct	acagttttat	cttttttagtg	360
tgcattgtgt	ctcctTTTTT	tttgcaata	gcttcacct	tataataact	catccatttt	420
attagtacat	ccatttaggg	tttaggggtta	atgggttttta	tagactaatt	tttttagtac	480
atctattttta	ttctattttta	gcctctaaat	taagaaaact	aaaactctat	tttagttttt	540
ttatttaata	atttagatat	aaaatagaat	aaaataaagt	gactaaaaat	taaacaata	600
ccctttaaga	aattaaaaaa	actaaggaaa	catttttctt	gtttcgagta	gataatgcc	660
gcctgttaa	cgccgtcgac	gagtctaacy	gacaccaacc	agcgaaccag	cagcgtcgcg	720
tccggccaag	cgaagcagac	ggcacggcat	ctctgtcgct	gcctctggac	ccctctcgag	780
agttccgctc	caccgttgga	cttgcctcgc	tgtcggcatc	cagaaattgc	gtggcggagc	840
ggcagacgtg	agccggcacg	gcaggcggcc	tcctcctcct	ctcacggcac	cggcagctac	900
gggggattcc	tttcccaccg	ctccttcgct	ttcccttcct	cggccgccgt	aataaataga	960
cacccctccc	acaccctctt	tccccaacct	cgtgtgtgtc	ggagcgcaca	cacacacaac	1020
cagatctccc	ccaaatccac	ccgtcggcac	ctccgcttca	aggtacgccg	ctcgtcctcc	1080
ccccccccc	ctctctacct	tctctagatc	ggcgttcggg	tccatgggtta	gggcccggta	1140
gttctacttc	tgttcatgtt	tgtgttagat	ccgtgtttgt	gttagatccg	tgctgctagc	1200
gttcgtacac	ggatgcgacc	tgtacgtcag	acacgttctg	attgctaact	tgccagtgtt	1260
tctctttggg	gaatcctggg	atggctctag	ccgttccgca	gacgggatcg	atttcatgat	1320
tttttttgtt	tcgttgcata	gggtttgggt	tgcccttttc	ctttatttca	atatatccg	1380
tgcacttggt	tgcgggtca	tcttttcagt	cttttttttg	tcttggttgt	gatgatgtgg	1440
tctggttggt	cggtcggtct	agatcggagt	agaattctgt	ttcaaaactac	ctggtggatt	1500
tattaatttt	ggatctgtat	gtgtgtgcca	tacatattca	tagttacgaa	ttgaagatga	1560
tggatggaaa	tatcgatcta	ggataggtat	acatgttgat	gcgggtttta	ctgatgcata	1620
tacagagatg	ctttttgttc	gcttggttgt	gatgatgtgg	tgtggttggg	cggtcgttca	1680
ttcgttctag	atcggagtag	aatactgttt	caaactacct	ggtgtattta	ttaattttgg	1740
aactgtatgt	gtgtgtcata	catcttcata	gttacgagtt	taagatggat	ggaaatatcg	1800
atctaggata	ggtatacatg	ttgatgtggg	ttttactgat	gcatatacat	gatggcatat	1860
gcagcatcta	ttcataatgct	ctaaccctga	gtacctatct	attataataa	acaagtatgt	1920
tttataatta	ttttgatctt	gatatacttg	gatgatggca	tatgcagcag	ctatatgtgg	1980
atTTTTTTTg	ccctgccttc	atacgtattt	tatttgcttg	gtactgtttc	ttttgtcgat	2040
gctcaccctg	ttgtttgggtg	ttacttctgc	agggatcccc	gatcatgcaa	aaactcatta	2100
actcagtga	aaactatgcc	tggggcagca	aaacggcggt	gactgaactt	tatggatagg	2160
aaaatccgtc	cagccagccg	atggccgagc	tgtggatggg	cgcacatccg	aaaagcagtt	2220
cacgagtga	gaatgccgcc	ggagatatcg	tttactgcg	tgatgtgatt	gagagtata	2280
aatcgactct	gctcggagag	cccggttgca	aacgctttgg	cgaactgcct	ttcctgttca	2340
aagtattatg	cgcagcacag	ccactctcca	ttcaggttca	tccaaacaaa	cacaattctg	2400
aaatcgggtt	tgccaaagaa	aatgccgcag	gtatcccgat	ggatgccgcc	gagcgttaact	2460
ataaagatcc	taaccacaag	cggagctgg	tttttgctct	gacgcctttc	cttgcatga	2520
acgcgtttcg	tgaattttcc	gagattgtct	ccctactcca	gccggctcgca	ggtgcacatc	2580
cggcgattgc	tcacttttta	caacagcctg	atgccgaacy	tttaagcgaa	ctgttcgcca	2640
gcctgttgaa	tatgcagggt	gaagaaaaat	cccgcgctct	ggcgatttta	aaatcggccc	2700
tcgatagcca	gcagggtgaa	ccgtggcaaa	cgattcgttt	aatttctgaa	ttttaccggy	2760
aagacagcgg	tctgttctcc	ccgtatttgc	tgaatgtgg	gaaattgaac	cctggcggaag	2820
cgatgttcc	gttcgctgaa	acaccgcacg	cttacctgca	aggcgtggcg	ctggaagtga	2880
tggcaaaact	cgataacgtg	ctgcgtgcgg	gtctgacgcc	taaatacatt	gatattccgg	2940
aactgggttc	caatgtgaaa	ttcgaagcca	aaccggctaa	ccagttgttg	accagcggg	3000
tgaacaagg	tgcagaactg	gacttcccga	ttccagtgg	tgattttgcc	ttctcgtctc	3060
atgaccttag	tgataaagaa	accaccatta	gccagcagag	tgcggccatt	ttgttctgcg	3120
tcgaaggcga	tgcaacgttg	tggaaagggt	ctcagcagtt	acagcttaaa	ccgggtgaat	3180
cagcgtttat	tgcggccaac	gaatcaccgg	tgactgtcaa	aggccacggc	cgttttagcgc	3240
gtgtttacaa	caagctgtaa	gagcttactg	aaaaaattaa	catctcttgc	taagctggga	3300
gctcgatccg	tcgacctgca	gatcgttcaa	acatttggca	ataaagtttc	ttaagattga	3360

atcctgttgc	cggctcttgcg	atgattatca	tataatttct	gttgaattac	gttaagcatg	3420
taataattaa	catgtaatgc	atgacgttat	ttatgagatg	ggtttttatg	attagagtcc	3480
cgcaattata	catttaatac	gcgataaaaa	acaaaatata	gcgcgcaaac	taggataaat	3540
tatcgcgcg	ggtgtcatct	atgttactag	atctgctagc	cctgcaggaa	atttaccggt	3600
gcccgggcgg	ccagcatggc	cgtatccgca	atgtgttatt	aagttgtcta	agcgtcaatt	3660
tgtttacacc	acaatatatc	ctgccaccag	ccagccaaca	gctccccgac	cggcagctcg	3720
gcacaaaatc	accactcgat	acaggcagcc	catcagaatt	aattctcatg	tttgacagct	3780
tatcatcgac	tgcacgggtc	accaatgctt	ctggcgctcag	gcagccatcg	gaagctgtgg	3840
tatggctgtg	caggctcgtaa	atcactgcat	aattcggtgc	gctcaaggcg	cactcccgtt	3900
ctggataatg	ttttttgctg	cgacatcata	acggttctgg	caaataattct	gaaatgagct	3960
gttgacaatt	aatcatccgg	ctogtataat	gtgtggaatt	gtgagcggat	aacaatttca	4020
cacaggaaac	agaccatgag	ggaagcgttg	atcgccgaag	tatcgactca	actatcagag	4080
gtagtgtggc	tcacgagcgc	ccatctcgaa	ccgacgttgc	tggccgtaca	tttgtacggc	4140
tccgcagtgg	atggcggcct	gaagccacac	agtgatattg	atttgctggt	tacggtgacc	4200
gtaaggcttg	atgaaacaac	gcggcgagct	ttgatcaacg	accttttgga	aacttcggct	4260
tccctgggag	agagcgagat	tctccgcgct	gtagaagtca	ccattgttgt	gcacgacgac	4320
atcattccgt	ggcgttatcc	agctaagcgc	gaactgcaat	ttggagaatg	gcagcgcaat	4380
gacattcttg	caggatatctt	cgagccagcc	acgatcgaca	ttgatctggc	tatcttgcgt	4440
acaaaagcaa	gagaacatag	cgttgccttg	gtaggtccag	cggcggagga	actctttgat	4500
ccggttcctg	aacaggatct	atttgaggcg	ctaaatgaaa	ccttaacgct	atggaactcg	4560
ccgcccgact	gggctggcga	tgagcgaat	gtagtgttta	cgttgtcccg	catttggtag	4620
agcgcagtaa	ccggcaaaat	cgcccggaag	gatgtcgctg	ccgactgggc	aatggagcgc	4680
ctgccggccc	agtatcagcc	cgtcatactt	gaagctaggc	aggcttatct	tggacaagaa	4740
gatcgcttgg	cctcgcgcg	agatcagttg	gaagaatttg	ttcactacgt	gaaaggcgag	4800
atcaccaagg	tatgcggcaa	ataaagctct	actggatctc	cgtacccccg	ggggatctgg	4860
ctcgccggcg	acgcacgacg	ccggggcgag	accataggcg	atctcctaaa	tcaatagtag	4920
ctgtaacctc	gaagcgtttc	acttgtaaca	acgattgaga	atttttgtca	taaaattgaa	4980
atacttgggt	cgcatttttg	tcacccgcgc	tcagccgcaa	ttctgacgaa	ctgcccattt	5040
agctggagat	gattgtacat	ccttcacgtg	aaaatttctc	aagcgtgtg	aacaagggtt	5100
cagatttttag	attgaaaggt	gagccgttga	aacacgttct	tcttgctgat	gacgacgtcg	5160
ctatgcggca	tcttattatt	gaatacctta	cgatccacgc	cttcaaagtg	accgcggtag	5220
ccgacagcac	ccagttcaca	agagtactct	cttcgcgcac	ggtcgatgtc	gtggtgtgtg	5280
atctaaattt	aggtcgtgaa	gatgggctcg	agatcgttcg	taatctggcg	gcaaagtctg	5340
atattccaat	cataattatc	agtggcgacc	gccttgagga	gacggataaa	gttgttgcac	5400
tcgagctagg	agcaagtgat	tttatcgcta	agccgttcag	tatcagagag	tttctagcac	5460
gcattcgggt	tgccttgccg	gtgcgcccc	acgttgtccg	ctccaaagac	cgacggtctt	5520
tttgttttac	tgactggaca	cttaatctca	ggcaacgtcg	cttgatgtcc	gaagctggcg	5580
gtgaggtgaa	acttacggca	ggtgagttca	atcttctcct	cgcgttttta	gagaaacccc	5640
gcgacgttct	atcgcgcgag	caacttctca	ttgccagtcg	agtacgcgac	gaggagggtt	5700
atgacaggag	tatagatgtt	ctcattttga	ggctgcgccc	caaacttgag	gcagatccgt	5760
caagccctca	actgataaaa	acagcaagag	gtgcccgtta	tttctttgac	gcggacgtgc	5820
aggtttcgca	cggggggacg	atggcagcct	gagccaattc	ccagatcccc	gaggaaatcg	5880
cgtgagcgg	cgcaaaccat	ccggcccgg	acaaatcggc	gcggcgctgg	gtgatgacct	5940
ggtggagaag	ttgaaggccg	cgcaggccgc	ccagcggcaa	cgcacgagag	cagaagcacg	6000
ccccggtgaa	tcgtggcaag	cggccgctga	tcgaatccgc	aaagaatccc	ggcaaccgcc	6060
ggcagccggt	gcgccgtcga	ttaggaagcc	gcccaggggc	gacgagcaac	cagatttttt	6120
cgttccgatg	ctctatgacg	tgggcacccg	cgatagtcgc	agcatcatgg	acgtggccgt	6180
tttccgtctg	tcgaagcgtg	accgacgagc	tggcgaggtg	atccgctacg	agcttccaga	6240
cgggcacgta	gaggtttccg	cagggccggc	cggcatggcc	agtgtgtggg	attacgacct	6300
ggtactgatg	gcggtttccc	atctaaccga	atccatgaac	cgataaccggg	aagggaaggg	6360
agacaagccc	ggccgcgtgt	tccgtccaca	cgttgcggac	gtactcaagt	tctgccggcg	6420
agccgatggc	ggaaagcaga	aagacgacct	ggtagaaacc	tgcatcgggt	taaaccacc	6480
gcacgttgcc	atgcagcgta	cgaagaaggc	caagaacggc	cgcttggtga	cggatatccga	6540
gggtgaagcc	ttgattagcc	gctacaagat	cgtaaagagc	gaaaccgggc	ggccggagta	6600
catcgagatc	gagctagctg	attggatgta	ccgcgagatc	acagaaggca	agaaccggga	6660

cgtgctgacg	gttcaccccc	attacttttt	gatcgatccc	ggcatcggcc	gttttctcta	6720
ccgcctggca	cgccgcgcgc	caggcaaggg	agaagccaga	tggttggtca	agacgatcta	6780
cgaacgcagt	ggcagcgccg	gagagttcaa	gaagttctgt	ttcaccgtgc	gcaagctgat	6840
cgggtcaaat	gacctgccgg	agtacgattt	gaaggaggag	gcggggcagg	ctggccccgat	6900
cctagtcagt	cgctaccgca	acctgatcga	gggcgaagca	tccgcgggtt	cctaattgtac	6960
ggagcagatg	ctagggcaaa	ttgccctagc	aggggaaaaa	ggtcgaaaag	gtctctttcc	7020
tgtggatagc	acgtacattg	ggaacccaaa	gccgtacatt	gggaaccgga	acccgtacat	7080
tgggaaccca	aagccgtaca	ttgggaaccg	gtcacacatg	taagtgactg	atataaaaaga	7140
gaaaaaaggc	gattttttccg	cctaaaactc	tttaaaactt	attaaaactc	ttaaaacccg	7200
cctggcctgt	gcataactgt	ctggccagcg	cacagccgaa	gagctgcaaa	aagcgcctac	7260
ccttcgggtcg	ctgcgtctcc	tacgccccgc	cgttcgcgtg	cggcctatcg	cggccgctgg	7320
ccgctcaaaa	atggctggcc	tacggccagg	caatctacca	gggcgcggac	aagccgcgcc	7380
gtcgccactc	gaccgcgggc	gctgaggtct	gcctcgtgaa	gaaggtgttg	ctgactcata	7440
ccaggcctga	atcgccccat	catccagcca	gaaagtgagg	gagccacggt	tgatgagagc	7500
tttggtgtag	gtggaccagt	tgggtgattt	gaacttttgc	tttgccacgg	aacggtctgc	7560
gttgctcgga	agatgcgtga	tctgatcctt	caactcagca	aaagttcgat	ttattcaaca	7620
aagccgcccgt	cccgtcaagt	cagcgtaatg	ctctgccagt	gttacaacca	attaaccaat	7680
tctgattaga	aaaactcatc	gagcatcaaa	tgaactgca	atttattcat	atcaggatta	7740
tcaataccat	atttttgaaa	aagccgtttc	tgtaatgaag	gagaaaactc	accgaggcag	7800
ttccatagga	tggcaagatc	ctggtatcgg	tctgcgattc	cgactcgctc	aacatcaata	7860
caacctatta	atttccctc	gtcaaaaata	aggttatcaa	gtgagaaatc	accatgagtg	7920
acgactgaat	ccggtgagaa	tggcaaaaagc	tctgcattaa	tgaatcggcc	aacgcgcggg	7980
gagaggcggg	ttgcgtattg	ggcgctcttc	tccttcctcg	ctcactgact	cgctgcgctc	8040
ggctgctcgg	cgcggcgag	cgggtatcac	tcactcaaa	gcggtaatat	ggttatccac	8100
agaatcaggg	gataacgcag	gaaagaacat	gtgagcaaaa	ggccagcaaa	aggccaggaa	8160
ccgtaaaaaag	gccgcgttgc	tggcggtttt	ccataggctc	cgccccctg	acgagcatca	8220
caaaaatcga	cgtcaagtc	agaggtggcg	aaacccgaca	ggactataaa	gataccaggc	8280
gtttccccc	ggaagctccc	tctgctgctc	tcctgttcgg	acctgcccgc	ttaccggata	8340
cctgtccgcc	tttctccctt	cgggaagcgt	ggcgctttct	catagctcac	gctgtaggta	8400
tctcagttcg	gtgtaggctg	ttcgctccaa	gctgggctgt	gtgcacgaac	ccccggttca	8460
gcccgaccgc	tgcgccttat	ccggttaacta	tctgtcttgag	tccaacccgg	taagacacga	8520
cttatcgcca	ctggcagcag	ccactggtaa	caggattagc	agagcgagg	atgtaggcgg	8580
tgctacagag	ttcttgaagt	ggtggcctaa	ctacggctac	actagaagaa	cagtatttgg	8640
tatctgcgct	ctgctgaagc	cagttacctt	cggaaaaaga	gttggtagct	cttgatccgg	8700
caaacaaacc	accgctggta	gcggtgggtt	ttttgtttgc	aagcagcaga	ttacgcgcag	8760
aaaaaaagga	tctcaagaag	atcctttgat	cttttctacg	gggtctgacg	ctcagtgga	8820
cgaaaactca	cgttaaggga	ttttggtcat	gagattatca	aaaaggatct	tcacctagat	8880
ccttttgatc	cggaaattaat	tctgtgtggt	ggcatgcaca	tacaaatgga	cgaacggata	8940
aaccttttca	cgccctttta	aatatccgat	tattctaata	aacgctcttt	tctcttaggt	9000
ttaccgcga	atatatcctg	tcaaacactg	atagtttaaa	ctgaaggcgg	gaaacgacaa	9060
tctgatcatg	agcggagaat	taaggagatc	acgttatgac	ccccgcgat	gacgcgggac	9120
aagccgtttt	acgtttggaa	ctgacagaac	cgcaacgctg	caggaaattgg	ccgcagcggc	9180
catttaaatc	aattgggcgc	gccgaattcg	agctcggtag	aagcttggcg	cgccggtacc	9240
caattcccgga	tctagtaaca	tagatgacac	cgcgcgcgat	aatttatcct	agtttgcgcg	9300
ctatattttg	ttttctatcg	cgtattaaat	gtataattgc	gggactctaa	tcataaaaaac	9360
ccatctcata	aataacgtca	tgcattacat	gttaattatt	acatgcttaa	cgtaattcaa	9420
cagaaattat	atgataatca	tgcgaagacc	ggcaacagga	ttcaatctta	agaaacttta	9480
ttgccaaatg	tttgaacgat	cggggaaatt	cggggatcta	attcccagg	ctgtagccga	9540
cgatggtgcg	ccaggagagt	tgttgattca	ttgtttgcct	ccctgctgcg	gtttttcacc	9600
gaagtccatg	ccagtccagc	gtttttgcag	cagaaaagcc	gccgacttcg	gtttgcggtc	9660
gcgagtgaag	atccctttct	tgttaccgcc	aacgcgcgaat	atgccttgcg	aggtcgcaaa	9720
atcggcgaaa	ttccatacct	gttcaccgac	gacggcgctg	acgcgatcaa	agacgcggtg	9780
atacatatcc	agccatgcac	actgatactc	ttcaactccac	atgtcggtgt	acattgagtg	9840
cagcccggct	aacgtatcca	cgccgtattc	ggtgatgata	atcggtgat	gcagtttctc	9900
ctgccaggcc	agaagttctt	tttccagtac	cttctctgcc	gtttccaaat	cgccgctttg	9960

gacataccat	ccgtaataac	ggttcaggca	cagcacatca	aagagatcgc	tgatggatc	10020
ggtgtgagcg	tgcgagaaca	ttacattgac	gcaggtgatc	ggacgcgtcg	ggtcagatct	10080
acgcgttgct	tccgccagtg	gcgcgaaata	ttcccgtgca	ccttgccggac	gggtatccgg	10140
ttcgttggca	atactccaca	tcaccacgct	tgggtgggtt	ttgtcacgcg	ctatcagctc	10200
tttaatcgcc	tgtaagtgcg	cttgctgagt	ttcccgttg	actgcctctt	cgctgtacag	10260
ttctttcggc	ttgltgccc	cttcgaaacc	aatgcctaaa	gagagggtta	agccgacagc	10320
agcagtttca	tcaatcacca	cgatgccatg	ttcatctgcc	cagtccagca	tctcttcagc	10380
gtaagggtta	tgcgaggtag	ggtaggagtt	ggccccaatc	cagtccatta	atgcgtggtc	10440
gtgcaccatc	agcacgttat	cgaatccttt	gccacgcaag	tccgcatctt	catgacgacc	10500
aaagccagta	aagtagaacg	gtttgtgggt	aatcaggaac	tggtcgcctt	tcactgccac	10560
tgaccggatg	ccgacgcgaa	gcgggtagat	atcacactct	gtctggcttt	tggctgtgac	10620
gcacagttca	tagagataac	cttcacccgg	ttgccagagg	tgccgattca	ccacttgcaa	10680
agtcccgtca	gtgccttgtc	cagttgcaac	cacctgttga	tccgcatcac	gcagttcaac	10740
gctgacatca	ccattggcca	ccacctgcca	gtcaacagac	gcgtgggttac	agtcttgccg	10800
gacatgcgtc	accacgggtga	tatcgctccac	ccaggtgttc	ggcgtgggtgt	agagcattac	10860
gctgcgatgg	attccggcat	agttaaagaa	atcatggaag	taagactgct	ttttcttgcc	10920
gtttctgctc	gtaatcacca	ttcccggcgg	gatagtctgc	cagttcagtt	cgttgttcac	10980
acaaacggtg	atacctgcac	atcaacaaat	tttggctcata	tattagaaaa	gttataaatt	11040
aaaatataca	cacttataaa	ctacagaaaa	gcaattgcta	tatactacat	tcttttatct	11100
tgaaaaaaat	atttgaaata	ttatattact	actaattaat	gataattatt	atataatat	11160
caaaggtaga	agcagaaact	tacgtacact	tttcccggca	ataacatacg	gcgtgacatc	11220
ggcttcaaat	ggcgtatagc	cgccctgatg	ctccatcact	tcctgattat	tgaccacac	11280
tttgccgttaa	tgagtgaccg	catcgaaacg	cagcacgata	cgctggcctg	cccaaccttt	11340
cggatataaag	acttcgcgct	gataccagac	gttgcccgc	taattacgaa	tatctgcac	11400
ggcgaaactga	tcgttaaaac	tgccctggc	agcaattgcc	cggtcttctt	gtaacgcgct	11460
ttcccaccaa	cgctgatcaa	ttccacagtt	ttcgcgatcc	agactgaatg	cccacaggcc	11520
gtcaggtttt	ttgatttcac	gggttgggg	ttctacagga	cggaccatgg	tcgacctcga	11580
atcaaccact	ttgtacaaga	aagctgggtt	gacaaattaa	gttgtcagtg	tgctgaagtt	11640
atataatcaag	ataggaagg	tggagatgtc	tgaagaaaga	gcatgtccag	gatgtccaat	11700
ttatagggga	cgatcgatca	tatcgatcga	cgggttgggg	tggtacaacg	tacgtacaga	11760
tggaacgaa	actcgatcga	tctaagcgtg	aaacatgcct	acacgtggga	acacttgatga	11820
cttatgttcc	gggaacgact	gtttgagaca	catttgagct	agtctacttt	acctcgaaac	11880
tgggcaacat	gtaactctaa	agtctaaact	aggagggttg	tccgtagtgt	aaatactagt	11940
gtgttatgtc	actttgagag	actttcgaca	attaggaata	tgaacctttt	tatttattta	12000
gttatgtgct	accaaataata	gaggctccca	tcagatggcc	caataagcca	aagaataagc	12060
caaattttga	attttcaaac	ttaattttga	gattgattct	gagataatct	caacgtagtt	12120
tttttcagta	ttggctttta	agttaaaaaa	aacacatata	taaaagtttt	acctacacat	12180
ttatttttat	tctctaataa	gocgttttgg	cttattagaa	aaaaattcaa	ccaatggggg	12240
ccagagaatg	ttttacttct	gatggtttcc	tggagggtat	atagaagtga	tggtcaatgc	12300
ttcttcattg	atgcttctaa	ttctttcgag	gattttcctg	gttgctgtag	ttagtaactt	12360
tgaatgtgat	tggataaacc	ttaaattcaa	tatgatgtga	tagagaaggt	ggttgatttc	12420
gagtatgate	ggcttgctct	ggatgtttgc	agctagtatt	cttgggtgtc	ttaatacaag	12480
tttttttttc	gtggttctac	gctaagcctc	gcaaaaagg	agagaggagc	agtgtgcaca	12540
agattttcct	tacataatcc	ggcactatgc	acttaattag	catctcccta	atattctgat	12600
catgtggtgt	ccctagatag	atataatggg	agagccatga	ccaagagaaa	ggaggttcga	12660
acatacagag	ttattagaac	atatagccat	acacagtga	gttcttcgga	aatccaaaag	12720
agatgaacta	cgactctaag	tctgacatat	aagatcattc	actagcaaga	gagaaaacac	12780
atgacacaat	aggagtagga	tggtacaagc	ctgctttt			12817



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/08085

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/53 C12N15/82 C12N9/02 A01H5/00 A01N65/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EMBL, WPI Data, EPO-Internal, SEQUENCE SEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>DATABASE EMBL 'Online!  ACCESSION NO: AJ270938,  2 August 2000 (2000-08-02)  DUDLER, R.: "Oryza sativa mRNA for  lipoxygenase (rci-1) gene"  XP002182835  the whole document  -&amp; SCHAFFRATH ULRICH ET AL:  "Characterization of RCI-1, a  chloroplastic rice lipoxygenase whose  synthesis is induced by chemical plant  resistance activators."  EUROPEAN JOURNAL OF BIOCHEMISTRY,  vol. 267, no. 19, October 2000 (2000-10),  pages 5935-5942, XP002182831</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	27-36, 39, 40



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*8\* document member of the same patent family

Date of the actual completion of the international search

15 November 2001

Date of mailing of the international search report

17/12/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl  
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 01 36464 A (DUVICK JONATHAN P ;PIONEER HI BRED INT (US); HU XU (US); LU GUIHUA) 25 May 2001 (2001-05-25) SEQ ID NO:5 ---	1
X	WO 99 66057 A (DRAPER JOHN ;KENTON PAUL (GB); BIOGEMMA UK LTD (GB); PAUL WYATT (G) 23 December 1999 (1999-12-23) the whole document ---	1,13,15, 16,18-25
X	ZUO J ET AL: "CHEMICAL-INDUCIBLE SYSTEMS FOR REGULATED EXPRESSION OF PLANT GENES" CURRENT OPINION IN BIOTECHNOLOGY, LONDON, GB, vol. 11, no. 2, April 2000 (2000-04), pages 146-151, XP001039646 ISSN: 0958-1669 the whole document ---	1,15,16, 18-25
X	JEPSON I ET AL: "CHEMICAL-INDUCIBLE GENE EXPRESSION SYSTEMS FOR PLANTS - A REVIEW" PESTICIDE SCIENCE, ELSEVIER APPLIED SCIENCE PUBLISHER. BARKING, GB, vol. 54, no. 4, December 1998 (1998-12), pages 360-367, XP000804296 ISSN: 0031-613X the whole document ---	1,15,16, 18-25
X	WO 93 21334 A (CADDICK MARK XAVIER ;ZENECA LTD (GB); GREENLAND ANDREW JAMES (GB);) 28 October 1993 (1993-10-28) page 45, line 1 - line 10 ---	1,15,16, 18-25
X	EP 0 388 186 A (DU PONT) 19 September 1990 (1990-09-19) the whole document ---	1,15,16, 18-25
X	BEAUDOIN NATHALIE ET AL: "Developmental regulation of two tomato lipoxygenase promoters in transgenic tobacco and tomato" PLANT MOLECULAR BIOLOGY, NIJHOFF PUBLISHERS, DORDRECHT, NL, vol. 33, no. 5, 1997, pages 835-846, XP002165965 ISSN: 0167-4412 the whole document ---	1,14-16, 20-25

-/--

## INTERNATIONAL SEARCH REPORT

in International Application No

PCT/EP 01/08085

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROUSTER JACQUES ET AL: "Identification of a methyl jasmonate-responsive region in the promoter of a lipoxygenase 1 gene expressed in barley grain." PLANT JOURNAL, vol. 11, no. 3, 1997, pages 513-523, XP002182954 ISSN: 0960-7412 the whole document	1,14-16, 20-25
X	SCHWEIZER PATRICK ET AL: "Different patterns of host genes are induced in rice by Pseudomonas syringae, a biological inducer of resistance, and the chemical inducer benzothiadiazole (BTH)." EUROPEAN JOURNAL OF PLANT PATHOLOGY, vol. 105, no. 7, October 1999 (1999-10), pages 659-665, XP001031411 ISSN: 0929-1873 page 661 -page 662 page 664	27-36
X	DATABASE EMBL 'Online! ACCESSION NO: AF002537, 1 July 2000 (2000-07-01) SASAKI, T., ET AL.: "Oryza sativa genomic DNA, chromosome 1, PAC clone:P0001B06." XP002182836 nts 97983-98231;	1,9,10, 13-16, 18-24
X	DATABASE EMBL 'Online! ACCESSION NO: AC073405, 28 June 2000 (2000-06-28) HSING, Y.- I.C., ET AL.: "Oryza sativa chromosome 5 clone P0036D10, complete sequence." XP002182837 nts 81014-81308	1,9,10, 13-16, 18-24
X	DATABASE EMBL 'Online! ACCESSION NO: AQ328986, 11 January 1999 (1999-01-11) WING, R.A., ET AL.: " nbxb0044M05f CUGI Rice BAC Library Oryza sativa genomic clone nbxb0044M05f,genomic survey sequence." XP002182838 the whole document	1,9-16, 18-24
X	WO 00 34479 A (TOMITA HIROFUMI ;HASHIMOTO HISAKO (JP); MITSUBISHI CHEM CORP (JP)) 15 June 2000 (2000-06-15) SEQ ID NO:1	1,9,10, 13-16, 18-24

-/--

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PENG YOU-LIANG ET AL: "A novel lipooxygenase from rice: Primary structure and specific expression upon incompatible infection with rice blast fungus." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 5, 1994, pages 3755-3761, XP002182832 ISSN: 0021-9258 the whole document ---	27-32
X	DATABASE GENESEQ 'Online! ACCESSION NO: AAQ68842, 18 May 1995 (1995-05-18) XP002182839 the whole document -& DATABASE GENESEQ 'Online! ACCESSION NO: AAR61136, 18 May 1995 (1995-05-18) XP002182840 abstract -& DATABASE WPI Section Ch, Week 199437 Derwent Publications Ltd., London, GB; Class C06, AN 1994-298802 XP002182841 & JP 06 225774 A (MITSUI GYOSAI SHOKUBUTSU BIO KENKYUSHO), 16 August 1994 (1994-08-16) abstract ---	27-32, 39-44
X	VOEROES KIRSTEN ET AL: "Characterization of a methyljasmonate-inducible lipooxygenase from barley (Hordeum vulgare cv. Salome) leaves." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 251, no. 1-2, January 1998 (1998-01), pages 36-44, XP002105414 ISSN: 0014-2956 the whole document ---	27-32, 39-44
Y	WO 97 26364 A (TEXAS A & M UNIVERSITY SYST) 24 July 1997 (1997-07-24) the whole document ---	35-38
Y	WO 97 26364 A (TEXAS A & M UNIVERSITY SYST) 24 July 1997 (1997-07-24) the whole document ---	35-38
A	HAUSE BETTINA ET AL: "A jasmonate-responsive lipooxygenase of barley leaves is induced by plant activators but not by pathogens." JOURNAL OF PLANT PHYSIOLOGY, vol. 154, no. 4, April 1999 (1999-04), pages 459-462, XP000913613 ISSN: 0176-1617 the whole document ---	1-16, 18-34

-/--

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SCHWEIZER PATRICK ET AL: "Induced systemic resistance in wounded rice plants."  PLANT JOURNAL,  vol. 14, no. 4, May 1998 (1998-05), pages 475-481, XP002183030  ISSN: 0960-7412  page 478, right-hand column, line 1 - line 10</p> <p>---</p>	1-16, 18-34
A	<p>US 5 804 693 A (KESSMANN HELMUT ET AL)  8 September 1998 (1998-09-08)  example 58B</p> <p>---</p>	1-16, 18-34
A	<p>SCHWEIZER PATRICK ET AL: "Gene-expression patterns and levels of jasmonic acid in rice treated with the resistance inducer 2,6-dichloroisonicotinic acid."  PLANT PHYSIOLOGY (ROCKVILLE),  vol. 115, no. 1, 1997, pages 61-70, XP002182834  ISSN: 0032-0889  page 64</p> <p>---</p>	1-16, 18-34
A	<p>SEGUCHI K ET AL: "ACTION MECHANISM OF N CYANOMETHYL-2-CHLOROISONICOTINAMIDE IN CONTROLLING RICE BLAST DISEASE"  JOURNAL OF PESTICIDE SCIENCE,  vol. 17, no. 2, 1992, pages 107-113, XP001031412  ISSN: 0385-1559  the whole document</p> <p>---</p>	1-16, 18-34
A	<p>DATABASE WPI  Section Ch, Week 199219  Derwent Publications Ltd., London, GB;  Class C06, AN 1992-156057  XP002182955  &amp; JP 04 094687 A (MITSUI GYOSAI SHOKU),  26 March 1992 (1992-03-26)  abstract</p> <p>-----</p>	

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.2

Present claim 1 and those parts of claims 13-25 referring to claim1, relate to a product/compound/method defined by reference to a desirable characteristic or property, namely chemical inducibility without wound or pathogen inducibility. The claims cover all products/compounds/methods having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such products/compounds/methods. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product/compound/method by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the products/compounds/methods concerning promoter molecules with demonstrated ability to express associated sequences in plant cells in the claimed manner, more specifically the search has been conducted on the basis of the sequence listings that allow said result to be achieved. The search has not been extended to promoters functioning in other hosts.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0136464	A	25-05-2001	AU	1486901 A	30-05-2001
			WO	0136464 A2	25-05-2001
WO 9966057	A	23-12-1999	AU	4519199 A	05-01-2000
			EP	1088090 A2	04-04-2001
			WO	9966057 A2	23-12-1999
WO 9321334	A	28-10-1993	AU	3901993 A	18-11-1993
			EP	0637339 A1	08-02-1995
			WO	9321334 A1	28-10-1993
EP 0388186	A	19-09-1990	AU	638448 B2	01-07-1993
			AU	5285490 A	22-10-1990
			EP	0388186 A1	19-09-1990
			EP	0463056 A1	02-01-1992
			HU	58802 A2	30-03-1992
			JP	3046063 B2	29-05-2000
			JP	4504054 T	23-07-1992
			NZ	232950 A	23-12-1992
			WO	9011361 A1	04-10-1990
			US	5608143 A	04-03-1997
			US	5364780 A	15-11-1994
			ZA	9002047 A	27-11-1991
WO 0034479	A	15-06-2000	JP	2000166570 A	20-06-2000
			AU	1797400 A	26-06-2000
			WO	0034479 A1	15-06-2000
WO 9726364	A	24-07-1997	US	5844121 A	01-12-1998
			AU	722435 B2	03-08-2000
			AU	1535597 A	11-08-1997
			EP	0874907 A1	04-11-1998
			WO	9726364 A1	24-07-1997
			US	5942661 A	24-08-1999
US 5804693	A	08-09-1998	US	5614395 A	25-03-1997
			AU	1249295 A	01-08-1995
			EP	0733117 A1	25-09-1996
			WO	9519443 A2	20-07-1995
			US	6262342 B1	17-07-2001
			US	5654414 A	05-08-1997
			US	5689044 A	18-11-1997
			US	5650505 A	22-07-1997
			US	5777200 A	07-07-1998
			US	5880328 A	09-03-1999
			US	5856154 A	05-01-1999
			US	5767369 A	16-06-1998
			US	5847258 A	08-12-1998
			US	5942662 A	24-08-1999
			US	5789214 A	04-08-1998
			AU	3108089 A	14-09-1989
			CZ	8901435 A3	15-12-1999
			DK	109889 A	09-09-1989
			EP	0332104 A2	13-09-1989
			FI	891054 A	09-09-1989
			HU	54417 A2	28-02-1991
			IL	89495 A	31-08-1995
			JP	2009377 A	12-01-1990

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5804693	A	KR 247622 B1	15-03-2000
		NO 890972 A	11-09-1989
		NZ 228234 A	25-09-1992
		PT 89915 A ,B	10-11-1989
		RU 2130491 C1	20-05-1999
		DD 283647 A5	17-10-1990
		KR 225352 B1	15-10-1999
		PL 162317 B1	30-09-1993
		ZA 8901726 A	29-11-1989
		AU 642865 B2	04-11-1993
		AU 5218390 A	27-09-1990
		CA 2012778 A1	24-09-1990
		EP 0392225 A2	17-10-1990
		HU 60770 A2	28-10-1992
		JP 3035783 A	15-02-1991
		KR 212691 B1	02-08-1999
		NZ 233053 A	26-10-1993
		ZA 9002250 A	28-11-1990
		AU 6557494 A	08-11-1994
		WO 9424295 A1	27-10-1994
JP 4094687	A	26-03-1992	NONE